

Asymptotic properties of a Nadaraya-Watson type estimator for regression functions of infinite order*

Seok Young Hong[†] Oliver Linton[‡]

University of Cambridge

April 22, 2016

Abstract

We consider a class of nonparametric time series regression models in which the regressor takes values in a sequence space. Technical challenges that hampered theoretical advances in these models include the lack of associated Lebesgue density and difficulties with regard to the choice of dependence structure in the autoregressive framework. We propose an infinite-dimensional Nadaraya-Watson type estimator, and investigate its asymptotic properties in detail under both static regressive and autoregressive contexts, aiming to answer the open questions left by Linton and Sancetta (2009). First we show pointwise consistency of the estimator under a set of mild regularity conditions. Furthermore, the asymptotic normality of the estimator is established, and then its uniform strong consistency is shown over a compact set of logarithmically increasing dimension with respect to α -mixing and near epoch dependent (NED) samples. We specify the explicit rates of convergence in terms of the Lambert W function, and show that the optimal rate is of logarithmic order, confirming the existence of the curse of infinite dimensionality.

Keywords: Functional Regression; Nadaraya-Watson estimator; Curse of infinite dimensionality; Near Epoch Dependence.

2000 Mathematics Subject Classification: Primary 62G05; Secondary 62G08

1 Introduction

Nonparametric modelling is a common method for analyzing time series. A major advantage of this approach is that the relationship between the explanatory variables

*Part of this paper was written while the first author visited Institut de Mathématiques, Toulouse in November 2013. We thank Philippe Vieu for his kind hospitality and many helpful advice. We also thank John Aston, Jeroen Dalderop, Paul Doukhan, Jiti Gao, Mikhail Lifshits, Alexei Onatski, and Richard Nickl for their helpful comments. Special thanks to Alexey Rudenko for providing an original Russian photocopy of Sytaya (1974), and Hyungjin Lee for translating the paper. Financial support from the European Research Council (ERC-2008AdG-NAMSEF) is gratefully acknowledged.

[†]Statistical Laboratory, Department of Pure Mathematics and Mathematical Statistics, University of Cambridge, United Kingdom, email: syh30@cam.ac.uk

[‡]Faculty of Economics, University of Cambridge, United Kingdom, e-mail: obl20@cam.ac.uk

under study, denoted by $X = (X_1, \dots, X_d)^\top$, and the response, say Y , can be modelled without assuming any restrictive parametric or linear structure; see for example Härdle (1990), Bosq (1996), or Fan and Yao (2003) for a comprehensive review. The cost of allowing for this extended flexibility is known as the *curse of dimensionality*; Stone (1980, 1982) showed that given a fixed measure of smoothness β allowed on the regression function the best achievable convergence rate (in minimax sense) $n^{-\beta/(2\beta+d)}$ deteriorates dramatically as the dimension/order d increases.

In a time series context it is often reasonable and advantageous to model the dependence upon the infinite past. For example, the AR(d) model with $d = \infty$ naturally extends those classical linear models, and enables the influence of *all past information* to be taken into account in the modelling procedure, thereby allowing for maximal flexibility in making statistical inference. It can also be very useful for several semi-parametric applications, and for testing the martingale hypothesis or the efficient market hypothesis in economics, where the conditional mean $E(Y_t|\mathcal{F}_{t-1})$ is the object of main interest. By employing suitable bandwidth adjustments, potential summability issues from the bias of the estimator can be resolved, and also, the influence of distant covariates can be suitably downweighted. Not restricting the number of conditioning variables also has an advantage of avoiding the statistician's *a priori* choice of the order d based on some order determination principles whose validity is subject to be questioned in practical situations. For these reasons, we are motivated to study a class of nonparametric time series regression models of infinite order that covers both static regression and autoregression cases.

Pagan and Ullah (1988) proposed studying the case where $d \rightarrow \infty$ in the context of an econometric analysis of risk models. Doukhan and Wintenberger (2008) studied the autoregressive model with $d = \infty$ under their notion of weak dependence, and showed the existence of a stationary solution, see also Wu (2011). Linton and Sancetta (2009) tackled the estimation problem in the context of an autoregressive model and established uniform almost sure consistency with respect to stationary ergodic sample observations. There is a vast literature on statistical research on functional data; typical examples include curves and images, both of which are infinite-dimensional in nature. Masry (2005) provided a rigorous treatment of nonparametric regression with dependent functional data in which X lies in a semi-metric space. Further, Mas (2012) derived the minimax rate of convergence for nonparametric estimation of the regression function on strictly independent and identically distributed regressors. Ferraty and Vieu (2006) detailed a number of extensions and overview of nonparametric approaches in functional statistics literature. Geenens (2011) gave an updated accessible summary on nonparametric functional regression, and introduced the term “curse of infinite dimensionality”, which reflects the difficulties that are expected in nonparametric estimation of infinite dimensional objects due to extreme sparsity. We discuss in the next section the difference between the functional data framework and our discrete time framework.

Whereas nonparametric regression problems for vectorial regressor have been exhaustively studied in the literature, statistical theories for their infinite-dimensional extensions have not been fully established due to inevitable technical challenges that hindered their widespread popularity. An obvious difficulty stems from the fact that the usual notion of density $p(\cdot)$ does not exist; since there is no σ -finite Lebesgue measure in infinite-dimensional spaces, the Lebesgue density (with respect to the infinite

product of probability measures) cannot be defined via the Radon-Nikodym theorem. Consequently, standard asymptotic arguments for kernel estimators are no longer valid, notably Bochner's lemma:

$$\begin{aligned} \frac{1}{h^d} E \left[\mathcal{K}^j \left(\frac{x - X}{h} \right) \right] &= \int \mathcal{K}^j(u) p(x - uh) \, du \\ &\rightarrow p(x) \|\mathcal{K}\|_j^j \quad \text{as } h \rightarrow 0 \end{aligned} \quad (1)$$

(under suitable regularity conditions; $j = 1, 2$), and classical limiting theories in non-parametric literature cannot be readily extended.

In this paper, we consider an infinite dimensional analogue of the classical nonparametric regression approach. We propose a local constant type estimator and investigate its asymptotic properties. In particular, we show both pointwise and uniform consistency of the estimator and establish its asymptotic normality under both static regression and autoregressive contexts with respect to α -mixing and near epoch dependent (NED) samples, respectively. Under some special conditions on the bandwidth, the explicit rate of convergence is given by specifying the small deviation probabilities, and the existence of the *curse of infinite dimensionality* is confirmed. The cases of finite or increasing dimensional regressors are covered as special cases of our framework.

2 Some Preliminaries

Consider the following regression model:

$$Y = m(X) + \varepsilon \quad (2)$$

where the regressor $X = (X_1, X_2, \dots)^\top$ is a random element taking values in some sequence space S , the response Y is a real-valued variable, and the stochastic error ε is such that $E(\varepsilon|X) = 0$ a.s. The objective is to estimate the Borel function $m(\cdot) = E(Y|X = \cdot)$ based on n random samples observed from a strictly stationary data generating process $\{(Y_t, X_t) \in \mathbb{R} \times S\}_{t \in \mathbb{Z}^+}$ having some pre-specified dependence structure (see section 2.1 below).

This framework is related to the recent advances in statistical research on functional data, see Ramsey and Silverman (2002) for general introduction and statistical applications. Recently, successful attempts have been made to adopt the notion of nonparametric inference on functional statistics literature, and fundamental theories have been well-established; Ferraty and Romain (2010) gives a comprehensive review. A major issue in this field of research lies in extending the statistical theories of $\mathbb{R}^d$ to those of function spaces. In this literature, the attention is usually on smooth functions that are approximated and reconstructed from finely discretised grids on some compact interval. In contrast, the setup in our model (2) can be viewed as looking at an infinite number of discrete observations. Such a difference is reflected by the fact that the observed data is taken to be a discrete process $X = (X_s)$ with unbounded $s \in \mathbb{Z}^+$ so that $S = \{f|f : \mathbb{N} \rightarrow \mathbb{R}\}$, rather than $X = (X(s))$ with $s \in [0, T]^k$ so that $S = \{f|f : [0, T]^k \subset \mathbb{R}^k \rightarrow \mathbb{R}\}$, e.g. curves if $k = 1$, images if $k \geq 2$. Despite isomorphism between separable Hilbert spaces, the discrete nature of our setting has several fundamental distinctive features that allow us to look further into many practical applications, a notable example of which is dependent autoregressive modelling.

An immediate consequence of the framework is that the tuning parameter can be imposed for each dimension, allowing one to control the marginal influence of the regressors. For instance when it is sensible to postulate that the influence of distant covariates is getting monotonically downweighted, one may set the marginal bandwidths to increase in lags so as to impose higher amount of smoothing. Depending on the nature of the regressor, S may be taken as the space of all infinite real sequences $\mathbb{R}^{\mathbb{N}} := \prod_{j=1}^{\infty} \mathbb{R}_j$ formed by taking Cartesian products of the reals, or various types of its linear subspaces such as ℓ_{∞} , ℓ_p , c . We propose to take $S = \mathbb{R}^{\mathbb{N}}$ so as to refrain from imposing any prior restrictions with regard to the choice of the regressor; for example, taking S to be the space of bounded sequences excludes the possibility of the regressors having infinite supports (e.g. Gaussian process).

2.1 Dependence structure and leading examples

A distinctive characteristic of time series data is temporal dependence between observations. As in the usual multivariate framework, we need some suitable assumptions on the dependence structure between the samples in order to derive asymptotic theories and to obtain convergence rates of the estimator. In nonparametric time series literature, Rosenblatt (1956)'s α -mixing has been the *de facto* standard choice due to being the weakest among the mixing-type asymptotic independence conditions. To name a few earlier works, pointwise and uniform consistency of the local constant estimator were shown by Roussas (1990) and Andrews (1995), respectively, and the asymptotic normality was established by Fan and Masry (1992). Mixing conditions have also been widely used in the context of dependent functional observations, see e.g. Ferraty et al. (2005), Masry (2005), and Delsol (2009).

DEFINITION 1. A stochastic process $\{Z_t\}_{t=1}^{\infty}$ defined on some probability space $(\Omega, \mathcal{F}, P)$ is called α -mixing or strongly mixing (cf. 'jointly' α -mixing if Z_t is $\mathbb{R}^{d>1}$ -valued) if

$$\alpha(r) := \sup_{A \in \mathcal{F}_{-\infty}^r, B \in \mathcal{F}_{r+q}^{\infty}} |P(A \cap B) - P(A)P(B)|$$

is asymptotically zero as $r \rightarrow \infty$, where $\mathcal{F}_a^b$ is the σ -algebra generated by $\{Z_t; a \leq t \leq b\}$. In particular, we say the process is algebraically (resp. exponentially) α -mixing if there exists some $c, k > 0$ such that $\alpha(k) \leq cr^{-k}$ (resp. if there exists some $c, \varsigma > 0$ such that $\alpha(r) \leq ce^{-\varsigma r}$).

The popularity of the mixing condition stems from the fact that its associated probability theories have been extensively developed, allowing simpler theoretical derivations, see for instance Doukhan (1994) or Rio (2000) for a comprehensive survey. However, several drawbacks have been pointed out in the literature. First, it is a rather strong technical condition that is hard to verify in practice. Moreover, it is well known that even some basic processes are not mixing. e.g. AR(1) with Bernoulli innovations.

The primary limitation of mixing conditions that we should be aware of in our paper arises from the choice of the framework upon which our theories are developed. Recall that given the sample observations $\{Y_t, X_t\}_{t=1}^n$ the object of estimation is the conditional mean $E(Y_t|\mathcal{F})$ where $\mathcal{F}$ represents some information set determined by

the nature of the conditioning variables. There are two leading cases: the first case is the static regression where the information set is taken to mean $\sigma(X_{jt}; j = 1, 2, \dots)$, the σ -algebra generated by marginal regressors, for example exogenous variables in econometric applications. The second case is the autoregression, where $X_{tj} = Y_{t-j}$, in which case $\mathcal{F} = \mathcal{F}_{t-1}$ represents $\sigma(Y_s; s \leq t-1)$, the σ -algebra generated by the sequence of lags of the responses $(Y_s)_{s \leq t-1}$. In practice, both types of conditioning information may be taken into account simultaneously.

In the static regression case the usual joint α -mixing condition can be imposed on the sample data $\{Y_t, X_t\}$ as in the multivariate framework; since marginal regressors are observed at the same time t : $X_t = (X_{1t}, X_{2t}, \dots)^\top$, assuming joint dependence does not require additional adjustments. Further, it will be shown later that in some cases asymptotic properties of the estimator can be invariant to the cross-dependence structure of X . Joint mixing implies both marginal component processes and any measurable function thereof are mixing¹ (with at most the same mixing rate $\alpha(\cdot)$). However in the autoregression setting, since the regressors are taken to be the lags of the response variable, measurable functions of X depend on the infinite past of the response, i.e. $\Psi(Y_{t-1}, Y_{t-2}, \dots)$ and hence are *not* mixing in general². Due to this reason an alternative set of dependence conditions is necessary to establish asymptotic theories for the second framework; we shall adopt the notion of near epoch dependence for the autoregressive setting and deal with two leading cases separately.

DEFINITION 2. A stochastic process $\{Z_t\}_{t=1}^\infty$ defined on some probability space $(\Omega, \mathcal{F}, P)$ is called near-epoch dependent or stable (in L_2 -norm) on a strictly stationary α -mixing process $\{\eta_t\}$ if the stability coefficients $v_2(m) := E|Z_t - Z_{t,(m)}|^2$ is asymptotically zero as $m \rightarrow \infty$, where $Z_{t,(m)} = \Psi_m(\eta_t, \dots, \eta_{t-m+1})$ for some Borel function $\Psi_m : \mathbb{R}^m \rightarrow \mathbb{R}$.

A process that is “near epoch” dependent on a mixing sequence is influenced primarily by the “recent past” of the mixing process and hence asymptotically resembles its dependence structure; see Lu (2001) for a detailed review. Following the usual convention, e.g. Bierens (1983), we shall take $\Psi_m(\eta_t, \dots, \eta_{t-m+1}) = E(Z_t | \eta_t, \dots, \eta_{t-m+1})$. We will show that under suitable conditions similar asymptotic theories can be derived for both regression and autoregression cases.

2.2 Local Weighting

In order to extend the standard multivariate nonparametric theories to infinite-dimension and to conduct nonparametric inference on the object of interest $m(\cdot)$, we first fix the notion of (i) local weighting and (ii) the measure of closeness between the objects. In the finite dimensional case, these are respectively done by the (multivariate) kernels $\mathcal{K} : (\mathbb{R}^d, \|\cdot\|_2) \rightarrow [0, \infty)$ which control the way the weights are given based on the distance, and the bandwidth h that regulates the proximity allowed between two objects in the smoothing procedure. The function $\mathcal{K}$ is called the product kernel or the spherical kernel depending on the way it is constructed from the univariate kernel

¹The converse is not necessarily true unless the marginal processes are independent to others, see Bradley (2005).

²Except for some very special cases; Davidson (1994, Theorem 14.9) gives a set of technical conditions under which a process with infinite (linear) temporal dependence is α -mixing.

$K(\cdot)$, i.e. $\mathcal{K}^P(u) = \prod_j^d K_j(u_j)$ or $\mathcal{K}^S(u) = K(\|u\|_2)$.

To account for the infinite-dimensional nature of our framework, we extend the latter approach to construct the weighting functions. We let

$$\mathcal{K}(u) := K(\|u\|) \quad (3)$$

where u is an element of a separable Banach space $(S^*, \|\cdot\|)$, and the univariate kernel K is a density function with positive support, for instance $K(\cdot) = \sqrt{2/\pi} \exp(-\cdot^2/2)$. We shall group the kernels into three subcategories depending on the way how they are generated. The first two, referred to as Type-I and Type-II kernels in Ferraty and Vieu (2006) generalize the usual ‘window’ kernels and monotonically decreasing kernels in finite dimension, respectively. Both types of kernels are continuous on compact support $[0, \lambda]$.

DEFINITION 3. *A function $K : [0, \infty) \rightarrow [0, \infty)$ is called a kernel of type-I if it integrates to 1, and if there exist strictly positive real constants C_1, C_2 (with $C_1 < C_2$) for which*

$$C_1 I_{[0, \lambda]}(u) \leq K(u) \leq C_2 I_{[0, \lambda]}(u), \quad (4)$$

where λ is some fixed positive real number. Also, a function $K : [0, \infty) \rightarrow [0, \infty)$ is called a kernel of type-II if it satisfies (4) with $C_1 \equiv 0$, and is continuous on $[0, \lambda]$ and differentiable on $(0, \lambda)$ with derivative K' that satisfies

$$C_3 \leq K'(u) \leq C_4$$

for some real constants C_3, C_4 such that $-\infty < C_3 < C_4 < 0$.

The definition above suggests that the uniform kernel on $[0, \lambda]$ is a type-I kernel, and the Epanechnikov, Biweight and Bartlett kernels belong to the class of Type-II kernels. Those with semi-infinite support, for example (one-sided) Gaussian, are covered by the last group, which we shall call the Type-III kernels.

DEFINITION 4. *A function $K : [0, \infty) \rightarrow [0, \infty)$ is a kernel of type-III if it integrates to 1, and if it is of exponential type; that is, $K(r) \propto \exp(Cr^\beta)$ for some constants β and C .*

2.3 Small deviation

The *small ball (or small deviation) probability* plays a crucial role in establishing the asymptotic theories in this paper. Let S^* be a separable sequence space that is complete with respect to some norm $\|\cdot\|$; then the small ball probability of an S^* -valued random element Z is a function defined as

$$\varphi_z(h) := P(\|z - Z\| \leq h). \quad (5)$$

The probability is called centered if $z = 0$, and shifted (with respect to a fixed point $z \in S^*$) if otherwise. The relation between the two quantities cannot be explicitly specified in general, and will be expressed in terms of the Radon-Nikodym derivative (See Assumption B1 below).

The name *small ball* stems from the fact that we are interested in its asymptotic behaviour as h , the bandwidth sequence in the context of nonparametric estimation, tends to zero. The function can be thought of as a measure for how much the observations are densely *packed* or *concentrated* around the fixed point z with respect to the associated norm and the reference distance h . From the definition it is straightforward to see that $\varphi(h) \rightarrow 0$ as $h \rightarrow 0$, and that $n\varphi(h)$ is an (approximate) count of the number of observations whose influence is taken into account in the smoothing procedure. When Z is a d -dimensional continuous random vector with Lebesgue density $p(\cdot) > 0$, it can be readily shown that the shifted small ball (with respect to the usual Euclidean norm) is given by

$$\varphi_z(h) = V_d h^d p(z) \propto h^d,$$

where $V_d = \pi^{d/2}/\Gamma(d/2 + 1)$ is the volume of d -dimensional unit sphere.

However when Z takes values in an infinite-dimensional space, it is generally difficult to specify the exact form of the small ball probability, and its behaviour varies depending heavily on the nature of the associated space and its topological structure. Due to non-equivalence of norms, it is intuitively clear that the *speed* at which $\varphi(h)$ converges to zero is affected by the choice of $\|\cdot\|$. In any case, a rapid decay is expected due to extreme sparsity of data in infinite dimensional spaces.

One possible example of S^* includes $(\ell_r, \|\cdot\|_r)$, the space of r -th power summable sequence equipped with the ℓ_r norm. As we shall reiterate below in the main text, we will take our main focus to be on the case of $r = 2$, since the behaviour of the small ball has been most studied in the case of square summable random variables. The case of $r > 2$ may also be derived without much difficulty, provided that the moment conditions are modified appropriately.

Writing the expected value of the kernel in terms of the small ball probability

$$\begin{aligned} EK\left(\frac{z - Z}{h}\right) &= EK\left(\frac{\|z - Z\|}{h}\right) \\ &= \int K(u) dP_{\|z - Z\|/h}(u) = \int K(u) d\varphi_z(uh), \end{aligned} \quad (6)$$

we are able to bypass the difficulties mentioned in the introduction, and establish the convergence of the integrals without explicitly requiring the existence of the density.

LEMMA 1. Ferraty and Vieu (2006, Lemma 4.4). *If K is a type-I kernel, then it satisfies*

$$C_1^j \leq \frac{1}{\varphi_z(h\lambda)} \int_0^\lambda K^j(v) d\varphi_z(vh) \leq C_2^j, \quad j = 1, 2 \quad (7)$$

where C_1, C_2 are strictly positive real constants defined in Definition 3. Furthermore, if the kernel K is type-II and if there exists some $\varepsilon_0 > 0$ and a constant $C_5 > 0$ such that $\forall \varepsilon < \varepsilon_0, \int_0^\varepsilon \varphi_x(u) du > C_5 \varepsilon \varphi_x(\varepsilon)$, then we have

$$C_6^j \leq \frac{1}{\varphi_z(h\lambda)} \int_0^\lambda K^j(v) d\varphi_z(vh) \leq C_7^j, \quad j = 1, 2 \quad (8)$$

where the constants $C_6 = -C_5 C_4$ and $C_7 = \sup_{s \in [0, \lambda]} K(s)$ are strictly positive.

Since Lemma 1 holds for every $h > 0$, it readily follows by the squeeze theorem that the following result holds for any kernels of type-I and II:

COROLLARY 1. *If the kernel K is either type-I or type-II, then for $j = 1, 2$ we have*

$$\frac{1}{\varphi_z(h\lambda)} E \left[\mathcal{K}^j \left(\frac{z - Z}{h} \right) \right] \rightarrow \xi_j \quad \text{as } h \rightarrow 0^+, \quad (9)$$

where ξ_1 and ξ_2 are some strictly positive real constants.

This result can be seen as an infinite-dimensional analogue of Bochner's lemma (1): i.e. for $Z \in \mathbb{R}^d$, $h^{-d} E \mathcal{K}((z - Z)/h) \rightarrow p(z) > 0$, and will play a fundamental role in constructing asymptotic theories of our estimator. It is obvious that ξ_j is bounded below and above by C_1^j and C_2^j , respectively (or C_6^j and C_7^j depending on the choice of the kernel). With specific choices of kernels and regressors the exact values of the constants may be specified.

REMARK. A natural question one may ask is whether (9) would still be valid for other kernels, in particular those with semi-infinite support, namely the Type-III kernels. In finite d -dimensional frameworks, it is well known that a set of standard assumptions including $\|u\|^d K(u) \rightarrow 0$ as $u \rightarrow \infty$ is sufficient for showing (1), see for instance Pagan and Ullah (1999, Lemma 1). However, in the infinite-dimensional setting the answer is negative in most usual cases where the kernel is of exponential type (e.g. Gaussian kernel). Whereas the lower bound of the limit can be easily constructed via Chebyshev's inequality:

$$(0 <) \exp(-c_\delta \delta) \leq [P(V \leq \delta)]^{-1} E \exp(-c_\delta V),$$

the upper bound may not exist, and the rate at which the small ball probability decays to zero may dominate the speed at which the integral (6) converges to zero. This claim cannot be formally verified for all general cases because (as mentioned above) there is no unified result for the asymptotic behaviour of small deviation available. Nonetheless, the idea can be sketched in the common case where the asymptotics of the distribution function (i.e. small deviation) is of exponential order: $P(V \leq \delta) \sim \exp(-C\delta^{-\theta/(\theta+1)})$ as $h \rightarrow 0$ for some constants C and θ ; by de Bruijn's exponential Tauberian theorem (see Bingham et al. (1987)) the necessary and sufficient condition for such a case is the following limiting behaviour of the Laplace transform near infinity:

$$E[\exp(-c_\delta V)] \sim \exp(C' \cdot c_\delta^{-\theta}) \quad \text{as } c_\delta \rightarrow \infty$$

for some constant C' . (e.g. Take, for the example of Gaussian kernel, $V = \|z - Z\|^2$, $\delta = h^2$, $c_\delta = 2^{-1}h^{-2}$) Difference in the order of convergence implies divergence of the limit (9). For this reason, we shall confine our attention to compactly supported kernels hereafter.

2.4 The Estimator

Consider the regression problem with $\mathbb{R}^N$ -dimensional regressor $X = (X_1, X_2, \dots)^\top$, where the regression operator is nonparametrically estimated with an estimator associated with a bandwidth matrix $H := \text{diag}(\underline{h}) = \text{diag}(h_1, h_2, \dots)$. No further condition

is explicitly imposed in the first instance except that the bandwidth sequence is appropriately chosen in such a way that a norm $\|\cdot\|$ can be admitted to the *weighted regressor*. As an expositional example, let us assume $h_j = \phi_j h$ where $\{\phi_j^{-1}\}_j$ is some square-summable sequence (e.g. $\phi_j = j^p$, for some $p \geq 1$).

Then by Kolmogorov's three-series theorem, the sequence of weighted regressor $\phi_j^{-1}X_j$ is square summable, w.p.1., provided that X'_j s are independent with finite variance and satisfy

$$\sum_{j=0}^{\infty} E \min \{1, \phi_j^{-2} X_j^2\} < \infty.$$

so that $(\phi_1^{-1}X_1, \phi_2^{-1}X_2, \dots)^\top =: Z$ is $(\ell_2, \|\cdot\|_2)$ -valued. Since this is a mild set of requirements, we see that sufficient flexibility is given while sophisticated mathematical developments are allowed. In the autoregressive context, the weighting sequence can be chosen to be non-decreasing so that the ‘‘relative influence’’ of the marginal regressor decreases in index (lag).

Seeing the arguments sketched above we shall assume from now on that Z is normed with $\|\cdot\| = \|\cdot\|_2$, and (with an abuse of notation) extend the usual definition of (shifted) small ball probability to account for varying bandwidths h_j and generalized support $[0, \lambda]$ as follows:

$$\varphi_x(\underline{h}\lambda) := P(\|H^{-1}(x - X)\| \leq \lambda), \quad (10)$$

or equivalently $P(X \in \mathcal{E}(x, \underline{h}\lambda))$, where $\mathcal{E}$ is the infinite-dimensional hyperellipsoid centred at $x \in \mathbb{R}^N$, and λ is the constant defined in section 2.2. Clearly, we have $\varphi_x(\underline{h}\lambda) = \varphi_z(h\lambda)$. For later reference, we also define the joint small ball probability as the joint distribution

$$\psi_x(\underline{h}\lambda; i, j) := P((X_i, X_j) \in \mathcal{E}(x, \lambda \underline{h}) \times \mathcal{E}(x, \lambda \underline{h})). \quad (11)$$

We are finally in a position to introduce our estimator. We propose to estimate $m(x) = E(Y|X = x)$, $x \in \mathbb{R}^N$ with the following local constant type estimator:

$$\widehat{m}(x) := \frac{\sum_{t=1}^n \mathcal{K}(H^{-1}(x - X_t)) Y_t}{\sum_{t=1}^n \mathcal{K}(H^{-1}(x - X_t))} \equiv \frac{\sum_{t=1}^n K(\|H^{-1}(x - X_t)\|) Y_t}{\sum_{t=1}^n K(\|H^{-1}(x - X_t)\|)}, \quad (12)$$

defined with respect to n -sample time series observations $\{Y_t, X_t\}_{t=1}^n$.

The estimator can be viewed as an infinite-dimensional generalization of the standard multivariate Nadaraya-Watson estimator, and is a special case of the one considered in Ferraty and Vieu (2002), Masry (2005) and references therein for general functional data. The aim of this paper is to examine some key asymptotic properties of the estimator and establish its statistical theory.

3 Asymptotic Properties

In this section we introduce the main results of our paper, deriving some large sample asymptotics of the proposed estimator (12). Consistency is shown in both pointwise and uniform sense and the asymptotic normality is established. Rates of convergence are specified under sets of regularity conditions. All proofs are detailed in the appendix.

We shall consider two different cases: 1) static regression and 2) autoregression. We therefore assume one of the two different sets of conditions on the dependence structure of the data generating process specified below. Assumption S1 below concerns with the static regression case where we have exogenous regressors that are jointly observed in time in a mildly dependent manner. No restriction is needed as regards cross-sectional dependence between the regressor, although necessary conditions will be imposed when needed at the later stage (see Assumptions D below). The second assumption concerns with the case where we observe a $\mathbb{R}$ -valued response that is α -mixing in time. Due to the reasons sketched in section 2.1, we adopt the notion of near epoch dependence to describe the dependence structure of infinite functions of the response variables. From the assumption we see that there is a trade-off between the degree of mixing rate and the possible order of moments allowed for the response variable $2 + \delta$.

ASSUMPTIONS S

- S1. *The marginal regressors $X_{1t}, X_{2t}, X_{3t}, \dots$ are exogenous variables, and the sample observations $\{Y_t, X_t\}_{t=1}^n = \{Y_t, (X_{1t}, X_{2t}, \dots)\}_{t=1}^n$ are (jointly) arithmetically α -mixing with rate $k > (2\delta + 4)/\delta$, where δ is as defined in Assumption A4 below.*
- S2. *The regressors are lags of the response, i.e. $X_{it} = Y_{t-i} \forall i$, and the process $\{Y_t\}_{t=1}^n$ is arithmetically α -mixing with $k > (2\delta + 4)/\delta$. Also, the process $K_t := K(\|H^{-1}(x - X_t)\|)$ is near epoch dependent on Y_t , and there exists some $m = m_n \rightarrow \infty$ such that the rate of stability for K_t denoted $v(m_n) = v(m)$ satisfies*

$$v(m)^{1/2}[\varphi_x(h\lambda)]^{-(2\delta+3)/(2\delta+2)-1}n^{1/(2(\delta+1))} \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (13)$$

3.1 Pointwise consistency

Pointwise consistency of the local constant estimator was first studied by Watson (1964) and Nadaraya (1964) with respect to i.i.d sample for the case of univariate regressor, i.e. $d = 1$. Their result was extended to the multivariate case by Greblicki and Krzyzak (1980) and Devroye (1981). Robinson (1983) and Bierens (1983) were amongst the earliest papers that worked on consistency of the estimator with respect to dependent data (both static regression and autoregression were allowed in their frameworks), followed by Roussas (1989), Fan (1990), and Phillips and Park (1998) to name a few out of numerous papers. The case of functional regressor was first studied by Ferraty and Vieu (2002), and references therein under various sets of regularity conditions.

In this section we show how these results can be extended specifically to our framework, proving pointwise weak consistency of the estimator (12) with respect to dependent sample satisfying either S1 or S2. A set of assumptions required for the theory is now introduced, and the series of arguments that leads to the conclusion is sketched.

ASSUMPTIONS A

- A1. *The regression operator $m : \mathbb{R}^N \rightarrow \mathbb{R}$ is continuous in some neighbourhood of x*

- A2. The marginal bandwidths satisfy $h_j = h_{j,n} \rightarrow 0$ as $n \rightarrow \infty$ for all $j = 1, 2, \dots$, where $\text{diag}(h_1, h_2, \dots) = \text{diag}(\underline{h}) = H$ is the bandwidth matrix, and the small ball probability obeys $n\varphi_x(\underline{h}\lambda) \rightarrow \infty$, where $\varphi_x(\underline{h}\lambda) := P(\|H^{-1}(x - X)\| \leq \lambda) \rightarrow 0$ as $n \rightarrow \infty$.
- A3. The kernel K is either type-I or type-II.
- A4. The response Y_t satisfies $E(|Y_t|^{2+\delta}) \leq C < \infty$ for some $C, \delta > 0$.
- A5. The joint small ball probability (11) satisfies $\psi_x(\underline{h}\lambda; i, j) = O(\varphi_x(\lambda\underline{h})^2)$, $\forall i \neq j$.
- A6. The conditional expectation $E(|Y_t Y_s| | (X_t, X_s)) \leq C < \infty$ for all t, s .

REMARK. Asymptotic unbiasedness of the estimator requires the continuity assumption A1; in other words, the estimator is unbiased at every point of continuity. Upon imposing further smoothness condition on the regression operator, the rate of convergence for the bias term can be specified (as we shall do later). Assumption A2 can be thought of as an extension of the usual bandwidth conditions that are assumed in finite-dimensional nonparametric literature. As discussed before, $n\varphi(\underline{h}\lambda)$ can be understood as an approximate number of observations that are “close enough” to x . Therefore, it is sensible to postulate that $n\varphi(\underline{h}\lambda) \rightarrow \infty$ as $n \rightarrow \infty$, meaning that the point x is visited many times by the sample of data as the size of sample grows to infinity. This is in line with the usual assumption $nh^d \rightarrow \infty$ when $X \in \mathbb{R}^d$, in which case the small ball probability is given by $\varphi(h) \propto h^d$. Condition A3 suggests that our result is valid under flexible choices of the compactly supported kernels. The standard moment condition A4 can be replaced by the boundedness condition $|Y| \leq C$, a.s. to simplify the proof, e.g. Billingsley’s inequality to bound asymptotic covariance terms. However we shall not consider this possibility here because this costs exclusion of the distributions with unbounded support. Conditions A5 and A6 are assumed to control the asymptotics of the covariance terms.

To sketch the idea, we write $K_t := K(\|H^{-1}(x - X_t)\|)$ for the sake of simplicity of presentation (note its dependence upon X_t), and express the estimator (12) as

$$\hat{m}(x) := \frac{\sum_{t=1}^n K(\|H^{-1}(x - X_t)\|) Y_t}{\sum_{t=1}^n K(\|H^{-1}(x - X_t)\|)} = \frac{\frac{1}{n} \sum_{t=1}^n \frac{K_t}{EK_1} Y_t}{\frac{1}{n} \sum_{i=1}^n \frac{K_i}{EK_1}} = \frac{\hat{m}_2(x)}{\hat{m}_1(x)}. \quad (14)$$

We then employ the following decomposition:

$$\begin{aligned} \hat{m}(x) - m(x) &= \frac{\hat{m}_2(x)}{\hat{m}_1(x)} - m(x) = \frac{\hat{m}_2(x) - m(x)\hat{m}_1(x)}{\hat{m}_1(x)} \\ &= \frac{E\hat{m}_2(x) - m(x)E\hat{m}_1(x)}{\hat{m}_1(x)} + \frac{[\hat{m}_2(x) - E\hat{m}_2(x)] - m(x)[\hat{m}_1(x) - E\hat{m}_1(x)]}{\hat{m}_1(x)}, \end{aligned} \quad (15)$$

where clearly $E\hat{m}_1(x) = 1$. We shall show consistency by proving that the ‘bias part’ $E\hat{m}_2(x) - m(x)$ and the ‘variance part’ $[\hat{m}_2(x) - E\hat{m}_2(x)] - m(x)[\hat{m}_1(x) - 1]$ are negligible in large sample. As for the latter term, it suffices to show mean squared convergence of $\hat{m}_2(x) - E\hat{m}_2(x)$ to zero because then $\hat{m}_1(x) \rightarrow^p 1$ will readily follow.

THEOREM 1. *Suppose A1-A5 hold. Then the estimator (12) with respect to the sample observations $\{Y_t, X_t^\top\}_{t=1}^n$ satisfying either S1 or S2 is weakly consistent for the regression operator $m(x)$. That is, as $n \rightarrow \infty$*

$$\widehat{m}(x) \xrightarrow{P} m(x). \quad (16)$$

REMARK. It is trivial to see that this result holds for i.i.d. data $\{(Y_t, X_t^\top); t \in \mathbb{Z}\}$ as well. The arguments simply becomes less involved because the covariance term does not need to be considered anymore. An alternative way to approach would be to apply a suitable exponential inequality, in which case some additional/different assumptions may be needed.

In the following section, we shall specify the convergence rates and establish the asymptotic normality by imposing a set of additional regularity conditions.

3.2 Asymptotic Normality

Rigorous studies on the limiting distribution of the standard Nadaraya-Watson estimator can be traced back to Schuster (1972) and Bierens (1987), where the case of univariate and multivariate regressors was considered, respectively. The case of dependent samples was studied by Robinson (1983), Bierens (1983) and then by many others under a wide range of different frameworks. Constructing asymptotic normality given a dependent sample data typically involves the use of Bernstein's "big-small" blocking argument followed by the standard Lindeberg-Feller central limit theorem.

General distributional theories for Nadaraya-Watson type estimators in a semi-metric space was established by Masry (2005, Theorem 4) and Delsol (2009). This paper is different from the existing literature in two perspectives. First, our specific setting allows autoregressive time series modelling with respect to dependent sample data. Second, whereas the final results of the existing papers were given in terms of abstract functions, our results are presented with the explicit rate of convergence, thereby allowing feasible applications.

The primary object of this section is to outline this procedure in detail, and to introduce the main theories and some interesting consequences thereof. To reiterate, both the i.i.d. case (in other words, when the marginal regressors X_j are independent and identically distributed) and also a dependent case shall be allowed. Specifically, we introduce how independence restriction can possibly be moderated to allow for a mild dependence structure. In particular, the second assumption below specifies the extent to which cross-sectional dependence can be allowed on the marginal regressors in our theory whilst allowing for specification of the exact form of the convergence rate of the estimator.

ASSUMPTIONS D. *The real-valued stochastic process $\{X_s\}_{s=1}^\infty$ satisfies $EX_s^4 \leq C < \infty$, and is either:*

D1. *independent and identically distributed, or*

D2. *stationary, and admits a causal moving average representation:*

$$X_s = \sum_{j=0}^{\infty} a_j \epsilon_{s-j}, \quad (17)$$

where a_j is a square summable sequence, and $\{\epsilon_j\}_j$ is an independent and identically distributed standard Gaussian sequence.

REMARK. In either case marginal regressors are required to be identically distributed; an additional distributional assumption is made in B2 below. Nonetheless, possible degree of dependence allowed in D2 is very mild and general. (In fact, the condition can even be generalized to require the existence of a non-causal MA representation, whose necessary and sufficient condition is simply the existence of the spectral density.) As an example, if a stationary stochastic process $\{X_s\}_{s=1}^\infty$ is α -mixing, then it always has a moving average representation of the form (17) provided it is Gaussian. This is because any α -mixing process is regular³ by definition, so is linearly regular when it is Gaussian, and hence (with stationarity) admitting the Wold decomposition with independent Gaussian innovations by Corollary 17.3.1 of Ibragimov and Linnik (1971). For this reason we see that each D1 and D2 is consistent with the case allowed in Assumption S1 and S2, respectively (because in Assumption S2 the regressors are cross-sectionally mixing as they are temporal lags of the response which is mixing by assumption), although the dependence structure specified in D2 can be allowed also for the static regression case (i.e. S1). This suggests that there is absolutely no need to assume independence between marginal regressors in our model (2) under Gaussianity. In particular, convergence rates of our estimator are shown to be invariant (upto some constant factor) to the choice between D1 and D2. Lastly, the requirement for finite fourth moment is to ensure that the squared marginal regressors have finite second moment (see below); obviously, in the autoregressive framework (Assumption S2), this forces δ in Assumption A4 to be greater than or equal to 2.

We now proceed to introduce some main assumptions needed for distributional theories.

3.2.1 The ‘bias component’

The first part concerns with the asymptotic ‘bias’, where Assumptions A is strengthened by imposing additional smoothness conditions and suitable bandwidth adjustments. It is a set of sufficient conditions under which one is able to specify the finite sample properties of the estimator/ the exact upper bound of the asymptotic bias. Note that alternatively, Fréchet differentiability type condition can be imposed, as was done in Mas (2012).

FURTHER ASSUMPTIONS A

A7. *Further to Assumption A2, the marginal bandwidths satisfy $h_j = \phi_j \cdot h$ for some positive real number ϕ_j where $h = h_n \rightarrow 0$ as $n \rightarrow \infty$.*

A8. *The regression operator $m : \mathbb{R}^N \rightarrow \mathbb{R}$ satisfies*

$$|m(x) - m(x')| \leq \sum_{j=1}^{\infty} c_j |x_j - x'_j|^\beta \quad (18)$$

³in the sense of e.g. Bulinskii and Shiryaev (2003, p.248)

for every $x', x = (x_1, x_2, \dots)^\top \in \mathbb{R}^\mathbb{N}$ and some real constant $\beta \in (0, 1]$, where c_j is some sequence of real constants for which $\sum_{j=1}^\infty c_j \leq 1$ and $\sum_{j=1}^\infty c_j \phi_j^\beta < \infty$.

REMARK. Roughly speaking, these assumptions are imposed to ensure the bias to be “well-behaved”. Assumption A7 extends the previous bandwidth condition A2. Obviously, it is consistent with what was previously assumed in A2 since $h \rightarrow 0$ implies coordinate-wise convergence of each marginal bandwidths in large sample. With this condition one is able to write the asymptotic bias expression and the order of the bias-variance balancing bandwidth in terms of the common factor h . It is possible to dispense with this condition at the cost of imposing minor modifications in A7; the asymptotic bias will then be written in terms of the infinite sum of a weighted marginal bandwidth h_j , whose convergence needs to be assumed. Furthermore, we remark that if the variance term is not concerned, the sequence of marginal coefficients ϕ_j does not necessarily have to be monotonically increasing in j , which would allow more flexible choices of bandwidths. However, we shall assume some increment condition in Assumptions B below, since such a restriction is needed for the variance component.

Assumption A8 replaces and strengthens Assumption A1, and can be thought of as a variant of the usual Hölder-type smoothness conditions. With the condition, the regression operator becomes a contraction mapping; it will be seen in the appendix that summability of the bias is ensured and the order of convergence rates for the bias term is specified, see also (20). One example of the contraction constant c_j would be $\exp(-j)$. Indeed, the contribution from each marginal dimension decrease in lag or index under this assumption.

In the context of autoregression where $X_j \equiv Y_{t-j}$ for any j , it is imperative to ensure the existence of stationary solution $\{Y_t\}$, since the model is essentially an infinite number of difference equations where finiteness of the solution is not automatically guaranteed. In the study of a class of general nonlinear AR(d) models, Dufflo (1997) and Götze and Hipp (1994) assumed what is called the Lipschitz mixing condition (or the strong contraction condition), which is essentially (18) replaced by finite d -sum on the right hand side. In our context, Assumption A6 plays such a role; Doukhan and Wintenberger (2008) showed that (18) with $\sum_{j=1}^\infty c_j < 1$, is sufficient for the existence of a stationary solution of the form

$$Y_t = f(\varepsilon_t, \varepsilon_{t-1}, \dots), \quad (19)$$

where f is some measurable function. Wu (2011) arrived at the same conclusion under the assumption of $\sum_{j=1}^\infty c_j = 1$; the restriction on c_j in our assumption reflects the findings of their papers.

3.2.2 The ‘variance component’

We now move on to the second chunk of assumptions that concerns with the ‘variance part’. As before, vectors Z and z are taken to mean $(\phi_1^{-1}X_1, \phi_2^{-1}X_2, \dots)^\top$ and $(\phi_1^{-1}x_1, \phi_2^{-1}x_2, \dots)^\top$, respectively, where the vector $x = (x_1, x_2, \dots)^\top$ is the point at which estimation is made.

ASSUMPTIONS B

- B1. *The induced probability measure P_{z-Z} is dominated by the measure P_Z , and its Radon-Nikodym density $dP_{z-Z}/dP_Z =: p^*$ is sufficiently smooth in some neighbourhood of zero, and is bounded away from zero at 0; i.e., $p^*(0) > 0$.*
- B2. *Given the regressor vector $X = (X_1, X_2, \dots)^\top \in \mathbb{R}^N$, the distribution F of X_s^2 is regularly varying near zero with strictly positive index.*
- B3. *Further to A5, the bandwidth sequence satisfies $h_j = j^p h$ with $p > 1$; i.e. $\phi_j = j^p$.*
- B4. *The conditional variance $\text{Var}[Y_t|X_t = u] = \sigma^2(u)$ is continuous in some neighbourhood of x ; i.e. $\sup_{u \in \mathcal{E}(x, h\lambda)} [\sigma^2(u) - \sigma^2(x)] = o(1)$. Similarly, the cross-conditional moment $E[(Y_t - m(x))(Y_s - m(x))|X_t = u, X_s = v] = \sigma(u, v)$, $t \neq s$ is continuous in some neighbourhood of (x, x) .*

REMARK. The first condition concerns with a transition of the shifted small ball probability to the centred quantity; it is taken from Mas (2012), where detailed discussions can be found. In the Gaussian framework, the limiting behaviour between the two is well-understood, see Zolotarev (1988) or de Acosta (1983) for example, unlike the non-Gaussian cases, see Skorohod (1967). Condition B2 is equivalent to saying

$$\lim_{x \rightarrow \infty} \frac{F(1/(\gamma x))}{F(1/x)} = \gamma^\rho$$

with strictly negative index of variation $\rho < 0$. This marks the close relationship between the laplace transform and the small ball probability, see Lifshits (1997), Mas (2012) for example for detailed discussions.

We require the function $F(1/x)$ to be regularly varying in order to ensure that the small ball probability *well-behaves* (near infinity) in asymptotic sense. Since only those having strictly negative ρ satisfy the condition, a regressor must have a distribution $F(1/x)$ that decreases (as $x \rightarrow \infty$) at a *reasonable speed*. By reasonable we mean that the relative weight of decrease follows a power law, and the variation should be continuous. A large class of common distributions satisfies the condition; for example, Gamma, Beta, Pareto, Exponential, Weibull, and also chi-squared distribution (in which case each X_s is Gaussian). We refer the reader to Conditions *I*, *L* and relevant discussions thereof in Dunker et al. (1998) for details. Next, the increment condition on the bandwidth B3 is assumed so as to specify the explicit behaviour of the small ball probability. We remark that different coefficients such as exponential weights (i.e. $h_j = e^j h$) can also be allowed, leading to similar distributional results given in the subsequent sections; however, we shall confine our attention to the case of polynomial law in this paper for the sake of simplicity of exposition. The standard conditions in B4 are assumed to deal with the asymptotics of the variance and covariance terms.

With reference to (15) we are now able to derive the following results for the bias

and variance components, in view of Assumptions A7, A8 and B, and Corollary 1:

$$\mathcal{B}_n := \left[E\widehat{m}_2(x) - m(x) \right] \leq h^\beta \lambda^\beta \sum_{j=1}^{\infty} c_j j^{p\beta} \quad (20)$$

$$\mathcal{V}_n := \text{Var} \left[\widehat{m}_2(x) - E\widehat{m}_2(x) \right] \simeq \frac{\sigma^2(x) \xi_2}{n \varphi_x(\underline{h} \lambda) \xi_1^2}, \quad (21)$$

where λ and $\widehat{m}_2(\cdot)$ are as in (4) and (14), respectively, and $a_n \simeq b_n$ means $a_n = b_n + o(1)$. With these results we are now ready to construct the asymptotic normality of our estimator.

3.2.3 Limiting distribution under independence

We first consider the situation in which there is a set of independent exogenous regressors in the static regression context; that is, when marginal regressors X_s are i.i.d., satisfying Assumption D1. In this case, the asymptotic normality can be established for a wide range of different choices of distributions.

Recall that under Assumption B2, the distribution function F of X^2 is regularly varying with the index of variation $\rho < 0$. Then by the characterisation theorem of Karamata (1933) (see for example Feller (1971)), there always exists a slowly varying function $\ell(x)$ that satisfies

$$F(1/x) = x^\rho \ell(x). \quad (22)$$

Now fix some p , the increment constant for bandwidth in Assumption B3, and denote by $\mathcal{L}(t)$ the Laplace transform of X^2 . Then we define the following constants

$$C_\ell = \lim_{h \rightarrow 0} \left[\ell^{-1/2} \left(h^{-\frac{4p}{2p-1}} \right) \right]$$

$$C^* = \frac{(2\pi)^{(1+2p\rho)} (2p-1)}{\Gamma^{-1}(1-\rho) \cdot (2p)^{\frac{2p(\rho+2)-1}{2p-1}}} \cdot \zeta^{\frac{2p(1+\rho)}{2p-1}}, \quad C^{**} = (2p-1) \cdot \left(\frac{K}{2p} \right)^{2p/(2p-1)}$$

and

$$\zeta = - \int_0^\infty \frac{u^{-1/2p} \mathcal{L}'(u)}{\mathcal{L}(u)} du, \quad \text{and} \quad \mathcal{V}_1 = \frac{C^* C_\ell \xi_2 \sigma^2(x)}{p^*(0) \xi_1^2 \lambda^{\frac{1+2p\rho}{2p-1}}},$$

where $\Gamma(\cdot)$ is the Gamma function, ξ_1 and ξ_2 are the constants specified in (9), λ is the upper bound of the support of the kernel, $p^*(\cdot)$ is the Radon-Nikodym derivative in B1. As pointed out by Dunker et al. (1999), the constant ζ may not exist in some cases.

To aid the exposition, we compute the constants for some common, regularly varying distributions in Table 1 below.

The main result of this subsection now follows. The theorem derives the limiting distribution of the infinite-dimensional Nadaraya-Watson type estimator for every i.i.d. regressor satisfying B2, and a set of sufficient regularity conditions under which the expression is valid. Notice that the specific choice of the regressor does not affect the rate of convergence.

THEOREM 2. *Suppose A2-A8 and B1-B4 hold. Let the marginal regressors X_s satisfy Assumption D1. Then the estimator (12) with respect to the sample observations*

$X_j^2 \sim F$ i.i.d.	ρ	$\lim_{x \rightarrow \infty} \ell(x) = C_\ell^{-2}$	ζ
Uniform(1,b)	-1	1	n/a
Gamma(α, β)	$-\alpha$	$\beta^\alpha \alpha^{-1} \Gamma(\alpha)^{-1}$	$\frac{\alpha \pi \beta^{-1/2p}}{\sin(\pi/2p)}$
$\exp(\eta)$	-1	η	$\frac{\pi \eta^{-1/2p}}{\sin(\pi/2p)}$
Weibull(α, β)	$-\alpha$	β	n/a
Pareto(θ, μ)	-1	μ/θ	n/a
χ_1^2	-1/2	$(2/\pi)^{1/2}$	$\frac{\pi 2^{(1-2p)/2p}}{\sin(\pi/2p)}$

Table 1: Examples of some key constants for common distributions

$\{Y_t, X_t^\top\}_{t=1}^n$ satisfying S1 is asymptotically normal with the following limiting distribution:

$$\sqrt{nh^{\frac{1+2\rho p}{2p-1}}} \exp \left\{ -C^{**}(\lambda h)^{-\frac{2}{2p-1}} \right\} \mathcal{V}_1^{-1} \left[\widehat{m}(x) - m(x) - \mathcal{B}_n \right] \implies N(0, 1), \quad (23)$$

where $\mathcal{B}_n = O(h^\beta)$ is the bias component as in (20).

3.2.4 Limiting distribution under gaussianity and dependence

The strict independence condition between the regressors assumed in the previous section can be relaxed to allow mild dependence specified in Assumption D2. In doing so, we shall make use of the arguments used in Hong, Lifshits and Nazarov (2015, Theorem 1), where the asymptotics of the small deviation probability is investigated under dependence. This setting not only grants sufficient flexibility in the static regression case, but moreover allows distributional result for the autoregressive context where the regressors constitute time lags of the response $X_s = Y_{t-s}$, with dependence structure as stipulated in Assumption S2 (and thereby satisfying D2). The cost required for this modification is Gaussianity assumption on the regressor.

With reference to Table 1, we can easily compute the constants C^* and C^{**} for the Gaussian case, denoted C_G^* and C_G^{**} respectively, as follows:

$$C_G^* = \frac{(2\pi)^{(1-p)}(2p-1)}{2 \cdot (2p)^{\frac{3p-1}{2p-1}}} \cdot \left[\frac{\pi 2^{(1-2p)/2p}}{\sin(\pi/2p)} \right]^{\frac{-p}{2p-1}}, \quad C_G^{**} = \frac{2p-1}{2} \left(\frac{\pi}{2p \sin \frac{\pi}{2p}} \right)^{\frac{2p}{2p-1}}.$$

With other constants defined as before, we now have the following asymptotic normality for the case of dependent regressors. We reiterate that the result covers both the regression and autoregression context (S1 and S2), and is invariant as long as the cross-sectional dependence structure satisfies Assumption D2.

For the square summable sequence a_j in (17) define

$$C_{\mathcal{A}} = \left[\frac{1}{2\pi} \int_0^{2\pi} \left| \sum_{j=0}^{\infty} a_j \exp(ijx) \right|^{1/p} dx \right]^p \quad \text{and} \quad \mathcal{V}_2 = \frac{C_G^* C_\ell \xi_2 \xi_1^{-2} \sigma^2(x)}{e^{-\frac{1}{2} \|\Gamma^{-1/2} z\|_2^2} (C_{\mathcal{A}} \lambda)^{\frac{1-p}{2p-1}}}.$$

where $\sigma^2(\cdot)$ is the conditional variance defined in Assumption B4 and $z = (z_j) = (j^{-p} x_j)$. Then we have the following result:

THEOREM 3. *Suppose A2-A8 and B1-B4 hold. Let the regressor $X = (X_1, X_2, \dots)^\top$ is jointly normally distributed with zero mean and the covariance operator Γ , and satisfies D2. Then, the estimator (12) with respect to sample observations $\{Y_t, X_t^\top\}_{t=1}^n$ satisfying either S1 or S2 is asymptotically normal with the following limiting distribution:*

$$\sqrt{nh^{\frac{1-p}{2p-1}} \exp \left\{ -C_G^{**} (C_A \lambda h)^{-\frac{2}{2p-1}} \right\} \mathcal{V}_2^{-1}} \left[\hat{m}(x) - m(x) - \mathcal{B}_n \right] \Rightarrow N(0, 1), \quad (24)$$

where $\mathcal{B}_n$ is again the bias component in (20).

REMARK. The additional term C_A is a function of the sequence a_j that reflects the dependence structure allowed between the regressors. This can be thought of as the “penalty term” for allowing dependence. The exponential term in the denominator of the asymptotic variance arises from the asymptotic equivalence relationship between the shifted and non-shifted small deviation for ℓ_2 -valued Gaussian variables, see (59) in the appendix.

3.3 Optimal Bandwidth

We now briefly discuss the issue of bandwidth choice. As in the finite dimensional framework, the results above confirm the existence of the trade-off relationship between the order of bias and variance. As the bandwidth grows, the variance decreases and the bound on the bias increases, vice versa. Therefore we may search for the optimal bandwidth h_{opt} that balances the order of those two quantities by solving their equivalence relation.

For example, as for the case of Gaussian regressor we have

$$h^\beta \sim \sqrt{\frac{\exp \left(Ch^{-2/(2p-1)} \right)}{nh^{\frac{1-p}{2p-1}}}} \quad (25)$$

so that

$$\left[2\beta + \frac{1-p}{2p-1} \right] \cdot \log h - Ch^{-\frac{2}{2p-1}} \sim -\log n.$$

Taking $h \sim (\log n)^a$ for some strictly negative a balances the leading terms on both sides:

$$\left[2\beta + \frac{1-p}{2p-1} \right] \cdot a \cdot \log \log n - (\log n)^{-\frac{2}{2p-1} \cdot a} \sim -\log n. \quad (26)$$

The explicit order a that solves (26) can be computed in terms of n , β and p . Writing $j := [2\beta + (1-p)/(2p-1)]$ and $k := 2/(2p-1)$ for the sake of simplicity, we can show after some computation that

$$a_{opt} = \frac{j \cdot \mathcal{W} \left(\frac{k}{j} \cdot n^{k/j} \right) - k \log n}{jk \cdot \log \log n}, \quad (27)$$

where $\mathcal{W}(y)$ is the lambert W function, which returns the solution x of $y = x \cdot e^x$. From this result the optimal bandwidth $h_{opt} \sim (\log n)^{a_{opt}}$ follows.

REMARK. We can search for the optimal bandwidth for the cases of non-Gaussian regressors by following exactly the same manner as above, so tedious details are omitted here. As regards the solution in (27), since the mapping $x \mapsto x \cdot e^x$ is not injective, the solution may be multi-valued on the negative domain, i.e. $y < 0$. This does not happen in (27) provided $\beta \geq 1/4$ (however big p is), because $(1-p)/(2p-1)$ is bounded away from $-1/2$.

Since the log terms dominate the double logarithm in (26) as the number of sample n increases, it can be readily expected that the optimal value of a in (27) converges to a limit in such a way that the leading orders are balanced. Below we introduce without formal justification a result that gives the lower bound (infimum) of the optimal bandwidth. Obviously, the result holds for any choice of the distribution of the regressor, since the order of the leading terms $-2/(2p-1)$ remains invariant as was shown in (23) and (24).

COROLLARY 2. *For any fixed choice of $p > 1$ and the distribution F of X^2 satisfying B2, the order of the optimal bandwidth a_{opt} satisfies*

$$a_{opt} \downarrow \left(-\frac{2p-1}{2} \right) \quad \text{as } n \rightarrow \infty, \quad (28)$$

which suggests that the lower bound of the optimal bandwidth is given by⁴

$$(\log n)^{-\frac{2p-1}{2}} \preceq h_{opt} \sim (\log n)^{a_{opt}}. \quad (29)$$

3.4 Uniform consistency

Uniform consistency of the Nadaraya-Watson estimator was first studied by Nadaraya (1964, 1970) and subsequently by numerous others. To name a few earlier literature, Devroye (1978) moderated the regularity conditions required in the previous papers, and Robinson (1983) proved consistency for dependent sample data. In functional statistics literature, only uniform consistency with respect to i.i.d. sample has been established so far, see Ferraty et al. (2010). Following their approach, we introduce the notion of Kolmogorov's entropy which accounts for the complexity of infinite dimensional spaces.

DEFINITION 5. *Given some $\eta > 0$, let $L(S, \eta)$ be the smallest number of open balls in E of radius η needed to cover the set $S \subset E$. Then Kolmogorov's η -entropy is defined as $\log L(S, \eta)$.*

From the definition it can be readily expected that Kolmogorov's entropy depends heavily on the nature/structure of the space we work on, and therefore is closely related to the rate of convergence of the estimator.

It is well known that the regression function cannot be estimated uniformly over the entire space because the magnitude of the regression function $m(\cdot)$ becomes non-predictable as $\|x\| \rightarrow \infty$, see Bosq (1996). In our framework, even greater restrictions

⁴The notation $f \preceq g$ means there exists some positive constant c such that $\lim_{n \rightarrow \infty} f(n)/g(n) \leq c$.

apply; since we are working on infinite sequence spaces, none of their subsets can be covered by a finite number of balls, and $L(S, \eta)$ then becomes infinite. Therefore, we propose to adopt an approximation argument and consider uniform consistency over a set whose effective dimension is increasing in sample size n . In particular, we define the set

$$S_\tau := \{u | (u_i)_{i \in \mathbb{Z}^+}, u_j = 0 \text{ for all } j > \tau, \|u\|_\infty \leq \lambda\} \quad (30)$$

where $\tau = \tau_n$ is some increasing sequence and fixed real number λ , and consider uniform consistency over this compact set. Kolmogorov's entropy of the set S_τ is given as follows:

LEMMA 2. *Kolmogorov's η -entropy of S_τ defined in (30) with $\tau = \tau_n (\rightarrow \infty)$ and $\lambda > 0$ is*

$$\log L(S_\tau, \eta) = \log \left[\left(\frac{2\lambda\sqrt{\tau}}{\eta} + 1 \right)^\tau \right]. \quad (31)$$

REMARK. We note that (31) is indeed in line with common intuition; as the dimension τ increases, the number of balls with some fixed radius required to cover the set goes off to infinity. The proof of this result can be done by exploiting the splitting technique and then by attempting to cover the polyhedron of increasing dimension. See appendix for details. From this result it readily follows that for fixed δ and λ , Kolmogorov's entropy $\log L(S_\tau, \eta)$ is of order $O(\tau + \tau \cdot \log \tau) = O(\tau \log \tau)$. As we shall discuss below, this has some implications with regard to the choice of the sequence τ_n .

We now introduce some further assumptions needed for uniform consistency:

ASSUMPTIONS C

C1. *The Kolmogorov's η -entropy $\log L(S_\tau, \eta)$ satisfies*

$$\frac{(\log n)^{3+\epsilon}}{n\varphi_x(\underline{h})} < \log L(S_\tau, \eta) < \frac{n\varphi_x(\underline{h})}{\log n} \quad \text{for some } \epsilon \in (0, 1) \quad (32)$$

Furthermore, $0 < \varphi_x(\underline{h}) \leq Ch < \infty$ and $(\log n)^2 / (n\varphi_x(\underline{h})) \rightarrow 0$ as $n \rightarrow \infty$.

C2. *The kernel function K is Lipschitz continuous on $[0, \lambda]$.*

REMARK. The first part of assumption C1 specifies the rate at which Kolmogorov's entropy should behave in sample size n (hence in dimension τ_n) in order for the theory to hold. From the upper and lower bound it readily follows that $n\varphi(h)$ must be of order strictly bigger than $(\log n)^{2+\epsilon/2}$. This assumption is general and is nothing restrictive; for example, taking $h \sim (\log n)^{-(2p-1)/2}$ in view of Corollary 2 and (27), it follows that $n\varphi(h) \sim (\log n)^{(2p-1)\beta}$. In this case, assumption (32) is valid as long as p is moderately large enough relative to $\beta \leq 1$. The second part of C1 is straightforward, and the last part only slightly strengthens the bandwidth condition in A2.

We now introduce the main result of this section. Note that with a slight abuse of notation X is hereafter taken to mean the regressor, but with zeros after its τ^{th} ($= \tau_n \rightarrow \infty$ as $n \rightarrow \infty$) entry; that is,

$$X = (X_1, X_2, \dots, X_\tau, 0, 0, \dots)^\top$$

(so that the original X is recovered as $n \rightarrow \infty$). Also, the regression operator and the estimator with respect to this truncated regressor are denoted by $m_\tau(\cdot)$ and $\hat{m}_\tau(\cdot)$, respectively.

A stonger condition is needed on the rate of mixing; from hereafter, by S1' and S2' we mean assumptions S1 and S2 but with the arithmetic mixing rate condition strengthened to exponential mixing (cf. Definition 1) with some exponent $\varsigma > 0$.

THEOREM 4. *Suppose that Assumptions A2-A8, B1-B3 and C1-C2 hold. Let the marginal regressors X_s satisfy D1, and let $\tau = \tau_n \sim (\log n)$. Then the estimator $\hat{m}_\tau(\cdot)$ with respect to sample observations $\{Y_t, X_t\}_{t=1}^n$ satisfying S1' is almost sure uniformly consistent for $m(x) = m(x_1, \dots)$ over S_τ :*

$$\sup_{x \in S_\tau} \left| \hat{m}_\tau(x) - m_\tau(x) \right| = O_{a.s.} \left(h^\beta + \sqrt{\frac{(\log n)^2 \exp \left\{ (h^2 + \tau^{1-2p})^{-1/(2p-1)} \right\}}{n(h^2 + \tau^{1-2p})^{\frac{1+2p\rho}{2(2p-1)}}}} \right), \quad (33)$$

If alternatively X_s is Gaussian and satisfies D2, then the same conclusion holds with respect to sample observations satisfying either S1' or S2':

$$\sup_{x \in S_\tau} \left| \hat{m}_\tau(x) - m_\tau(x) \right| = O_{a.s.} \left(h^\beta + \sqrt{\frac{(\log n)^2 \exp \left\{ (h^2 + \tau^{1-2p})^{-1/(2p-1)} \right\}}{n(h^2 + \tau^{1-2p})^{\frac{1-p}{2(2p-1)}}}} \right). \quad (34)$$

REMARK. The τ terms in the convergence rate reflects the penalty for truncation. Since $\tau \sim (\log n)$ the order of the term h^2 dominates that of τ^{1-2p} as long as $h \succeq (\log n)^{-(2p-1)/2}$. This can be done by choosing the bias-variance optimal bandwidth; following the same arguments in the pointwise case, choosing $h \sim (\log n)^a$ and solving for n we end up with

$$a_{opt} = \frac{j \cdot \mathcal{W} \left[\frac{k}{j} \exp(-\frac{k}{j} 2 \log \log n - k \log n) \right] + 2k \log \log n - k \log n}{jk \log \log n}.$$

And because the order of the leading terms is $(\log n)^{-(2p-1)/2}$ as in the pointwise case, it is straightforward to see that the lower bound of the optimal bandwidth in Corollary 2 still continues to hold; that is, $h_{opt} \succeq (\log n)^{-(2p-1)/2}$. This is again invariant to the choice of distribution F of the regressor.

We end up with the result that follows. Note that β is the measure of smoothness defined in assumption A8.

COROLLARY 3. *Suppose conditions assumed in Theorem 4 hold. Then, upon choosing $h \sim (\log n)^{a_{opt}}$, we have*

$$\sup_{x \in S_\tau} \left| \hat{m}_\tau(x) - m_\tau(x) \right| = O_{a.s.} \left([\log n]^{\beta \cdot a_{opt}} \right). \quad (35)$$

4 Appendix: Proofs of the main results

Throughout, C (or C' , C'') is taken to mean some generic constant that may take different values in different places unless defined specifically otherwise.

4.1 Proof of Theorem 1

Recalling the decomposition (15):

$$\widehat{m}(x) - m(x) = \frac{E\widehat{m}_2(x) - m(x)}{\widehat{m}_1(x)} + \frac{\widehat{m}_2(x) - E\widehat{m}_2(x)}{\widehat{m}_1(x)} - \frac{m(x)[\widehat{m}_1(x) - 1]}{\widehat{m}_1(x)},$$

we shall show $E\widehat{m}_2(x) - m(x) \rightarrow 0$ and $\widehat{m}_2(x) - E\widehat{m}_2(x) \rightarrow^p 0$, since $\widehat{m}_1(x) \rightarrow^p 1$ will then follow from the latter and complete the proof.

As regards the first ‘bias component’, denoting by $\mathcal{E}(x, \lambda \underline{h})$ the infinite dimensional hyperellipsoid centred at $x \in \mathbb{R}^N$ with semi-axes h_j in each direction we have

$$\begin{aligned} E\widehat{m}_2(x) - m(x) &= E\left(\frac{1}{nEK_1} \sum_{t=1}^n K_t Y_t - m(x)\right) \\ &= \frac{1}{EK_1} EK_1 Y_1 - \frac{EK_1}{EK_1} m(x) = \frac{1}{EK_1} E\left[E\left[(Y_1 - m(x))K_1 \middle| X\right]\right] \\ &= \frac{1}{EK_1} E\left[\left[m(X) - m(x)\right]K_1\right] \leq \sup_{u \in \mathcal{E}(x, \lambda \underline{h})} |m(u) - m(x)| \rightarrow 0 \end{aligned} \quad (36)$$

as $n \rightarrow \infty$, where K_t is the shorthand notation for $K(\|H^{-1}(x - X_t)\|)$ as introduced in the main text above. The second equality is justified by stationarity that is preserved under measurable transformation, and the last inequality is due to compact support of the kernel and continuity of the regression operator at x (Assumption A1).

The next step concerns with the ‘variance component’, where the mean-squared convergence of $\widehat{m}_2 - E\widehat{m}_2$ to zero will be shown. We write

$$\widehat{m}_2 - E\widehat{m}_2 = \frac{1}{n} \sum_{t=1}^n \frac{1}{EK_1} \left\{ K_t Y_t - E(K_t Y_t) \right\} =: \frac{1}{n} \sum_{t=1}^n Q_t. \quad (37)$$

The arguments to follow depend on the temporal dependence structure of Q_t . In the static regression case, Q_t is a measurable function of $Y_t, X_{1t}, X_{2t}, \dots$, and hence inherits their joint dependence structure. That is, Q_t is also arithmetically α -mixing with the rate specified in S1. In the autoregression case, we assume near epoch dependence of K_t (and hence Q_t ; upon assuming a slightly stronger condition on the stability coefficient) on Y_t as specified in Assumption S2, due to its dependence upon infinite past of Y_t . We shall hence deal with two cases separately.

CASE 1: STATIC REGRESSION. It is clearly sufficient to prove $\text{Var}(\widehat{m}_2 - E\widehat{m}_2) \rightarrow 0$ to show mean squared convergence. Since Q_t is (temporally) stationary we have

$$\text{Var}(\widehat{m}_2 - E\widehat{m}_2) = \frac{1}{n^2} \sum_{t=1}^n \text{Var}(Q_t) + \frac{2}{n^2} \sum_{1 \leq i < j \leq n} \text{Cov}(Q_i, Q_j) \quad (38)$$

$$\begin{aligned} &= \frac{1}{n} \text{Var}(Q_1) + \frac{2}{n^2} \sum_{1 \leq |i-j| < n} \text{Cov}(Q_i, Q_j) \\ &= \frac{1}{n} \text{Var}(Q_1) + \frac{2}{n^2} \sum_{s=1}^{n-1} (n-s) \cdot \text{Cov}(Q_1, Q_{s+1}) \\ &=: A_1 + A_2. \end{aligned} \quad (39)$$

Now, by (7), (8) and Assumption A it follows that

$$\begin{aligned} A_1 &= \frac{1}{nE^2K_1} \text{Var} \left(K_1 Y_1 - EY_1 K_1 \right) = \frac{\text{Var} (K_1 Y_1)}{nE^2K_1} \leq \frac{EK_1^2 Y_1^2}{nE^2K_1} \\ &= \frac{EK_1^2 Y_1^2 1\{|Y| \leq v\}}{nE^2K_1} + \frac{EK_1^2 Y_1^2 1\{|Y| > v\}}{nE^2K_1} \leq \frac{Cv^2}{n\varphi_x(\lambda \underline{h})} + \frac{CE|Y|^{2+\delta} v^{-\delta}}{n\varphi_x(\lambda \underline{h})}, \end{aligned} \quad (40)$$

where $v = v_n$ is some increasing truncation sequence. Then (40) tends to zero if the sequence is chosen to increase sufficiently slowly in such a way that $v = o(\sqrt{n\varphi})$.

We now move on to the second term A_2 and investigate the covariance term. Since measurable transformations of mixing variables preserve the mixing property, using Davydov's inequality, see Davydov (1968, Lemma 2.1) or Bosq (1996, Corollary 1.1), and boundedness of the kernel functions (4) we have

$$\begin{aligned} |\text{Cov}(Q_1, Q_{s+1})| &= \left| \text{Cov} \left(Y_1 \frac{K_1}{EK_1}, Y_{s+1} \frac{K_{s+1}}{EK_1} \right) \right| \\ &\leq \frac{12C \cdot \{E|Y_t|^{2+\delta}\}^{\frac{2}{2+\delta}} s^{-k\delta/(2+\delta)}}{\varphi_x(\underline{h}\lambda)^2}. \end{aligned} \quad (41)$$

In the meantime,

$$\begin{aligned} |\text{Cov}(Q_1, Q_{s+1})| &= \left| \text{Cov} \left(Y_1 \frac{K_1}{EK_1}, Y_{s+1} \frac{K_{s+1}}{EK_1} \right) \right| \\ &\leq \left| E \left(Y_1 \frac{K_1}{EK_1} Y_{s+1} \frac{K_{s+1}}{EK_1} \right) \right| + \left| E \left(Y_1 \frac{K_1}{EK_1} \right) E \left(Y_{s+1} \frac{K_{s+1}}{EK_1} \right) \right| \\ &\leq \frac{C}{\varphi_x(\underline{h}\lambda)^2} |E(K_1 K_{s+1})| + \frac{C}{E^2 K_1} |E(K_1) E(K_{s+1})| \\ &\leq \frac{C}{\varphi_x(\underline{h}\lambda)^2} \cdot \psi_x(\lambda \underline{h}; 1, s+1) + C'' \leq C'' \end{aligned} \quad (42)$$

by stationarity, law of iterated expectation, boundedness of regression function, and Assumption A6, A5 (along with the upper bound $\psi(\lambda \underline{h}; 1, s+1)$ of $EK_1 K_{s+1}$ obtained as a direct consequence of A5 following similar arguments used for Lemma 1).

With reference to (41) and (42), we take some increasing sequence $r_n \rightarrow \infty$ such that $r_n = o(n)$, and write

$$\begin{aligned} \sum_{s=1}^{n-1} |\text{Cov}(Q_1, Q_{s+1})| &= \sum_{s=1}^{r_n-1} |\text{Cov}(Q_1, Q_{s+1})| + \sum_{s=r_n}^{n-1} |\text{Cov}(Q_1, Q_{s+1})| \\ &\leq C''(r_n - 1) + \sum_{s=r_n}^{n-1} \frac{Cs^{-k\delta/(2+\delta)}}{\varphi_x(\underline{h}\lambda)^2} = O \left(r_n + \frac{r_n^{-k\delta/(2+\delta)+1}}{\varphi_x(\underline{h}\lambda)^2} \right), \end{aligned} \quad (43)$$

which is $O(\varphi_x(\underline{h}\lambda)^{-2(2+\delta)/(k\delta)})$ upon choosing $r_n \sim \varphi_x(\underline{h}\lambda)^{-2(2+\delta)/(k\delta)}$.

Consequently, since $k \geq 2(2 + \delta)/\delta$ it follows that

$$\begin{aligned}
A_2 &:= \frac{2}{n^2} \sum_{s=1}^{n-1} (n-s) \cdot \text{Cov}(Q_1, Q_{s+1}) = \frac{2}{n} \sum_{s=1}^{n-1} \left(1 - \frac{s}{n}\right) \cdot \text{Cov}(Q_1, Q_{s+1}) \\
&= O\left(\frac{1}{n[\varphi_x(\underline{h}\lambda)]^{2(2+\delta)/(k\delta)}} + \frac{1}{n^2[\varphi_x(\underline{h}\lambda)]^{2(2+\delta)/(k\delta)}}\right) \\
&= O\left(\frac{1}{n[\varphi_x(\underline{h}\lambda)]^{2(2+\delta)/(k\delta)}}\right) \rightarrow 0 \quad \text{as } n \rightarrow \infty
\end{aligned} \tag{44}$$

and the desired result is obtained.

CASE 2: AUTOREGRESSION. We return back to (37):

$$\widehat{m}_2 - E\widehat{m}_2 = \frac{1}{n} \sum_{t=1}^n \frac{1}{EK_1} \left\{ K_t Y_t - E(K_t Y_t) \right\} =: \frac{1}{n} \sum_{t=1}^n Q_t. \tag{45}$$

In this framework $K_t = K(\|H^{-1}(x - X_t)\|)$ is a function of $(Y_{t-1}, Y_{t-2}, \dots)$; despite loosing the mixing property, it inherits stationarity of the mixing process $\{Y_t\}$. We write $K_{t,(m)} = \Psi_m(Y_t, Y_{t-1}, Y_{t-2}, \dots, Y_{t-m+1}) = E(K_t | Y_t, \dots, Y_{t-m+1})$, where m is as in assumption S2. Clearly, $K_{t,(m)}$ preserves the mixing dependence structure of Y_t by elementary arguments, see e.g. Lu (2001, Remark 1(b)). Now write

$$\begin{aligned}
\widehat{m}_2 - E\widehat{m}_2 &= \frac{1}{n} \sum_{t=1}^n \frac{1}{EK_1} \left[K_{t,(m)} Y_t - E(K_{t,(m)} Y_t) \right] + \frac{1}{n} \sum_{t=1}^n \frac{1}{EK_1} \left[K_t Y_t - K_{t,(m)} Y_t \right] \\
&\quad + \frac{1}{n} \sum_{t=1}^n \frac{1}{EK_1} \left[E(K_{t,(m)} Y_t) - E(K_t Y_t) \right] = R_1 + R_2 + R_3,
\end{aligned} \tag{46}$$

and first consider the last term R_3 .

Fix some increasing sequence $q = q_n \rightarrow \infty$, and write $Y_{t,L} := Y_t 1_{\{|Y_t| \leq q\}}$ and $Y_{t,U} = Y_t 1_{\{|Y_t| > q\}}$. Then

$$\begin{aligned}
EY_t K_{t,(m)} &= EY_t K(\|H^{-1}(x - X_t)\|) - EY_{t,U} K(\|H^{-1}(x - X_t)\|) \\
&\quad + EY_{t,L} K_{t,(m)} - EY_{t,L} K(\|H^{-1}(x - X_t)\|) \\
&\quad + EY_{t,U} K_{t,(m)} \\
&= D_1 + D_2 + D_3.
\end{aligned} \tag{47}$$

The second part of D_1 is given by

$$\begin{aligned}
EY_{t,U} K(\|H^{-1}(x - X_t)\|) &\leq E|Y_t| 1_{\{|Y_t| > q\}} K(\|H^{-1}(x - X_t)\|) \\
&\leq q^{-(\delta+1)} E|Y_t|^{2+\delta} 1_{\{|Y_t| > q\}} K_t \\
&\leq Cq^{-(\delta+1)} E|Y_t|^{2+\delta} 1_{\{|Y_t| > q\}} = o(q^{-(\delta+1)})
\end{aligned} \tag{48}$$

because $1_{\{|Y_t| > q\}} = o(1)$ as $n \rightarrow \infty$. Following similar arguments on D_3 we have $D_1 + D_3 = EY_t K_t + o(q^{-(\delta+1)})$. So we are now left with the middle term D_2 :

$$D_2 \leq E|Y_{t,L}| |K_t - K_{t,(m)}| = O\left(q\sqrt{v_2(m_n)}\right) \tag{49}$$

by Hölder's inequality. Therefore in view of (47),(48) and (49) we have

$$R_3 = \frac{1}{nEK_1} \sum_{t=1}^n \left[EK_{t,(m)} Y_t - E(K_t Y_t) \right] = o\left(\frac{q^{-(\delta+1)}}{\varphi_x(\lambda \underline{h})}\right) + O\left(\frac{q\sqrt{v_2(m_n)}}{\varphi_x(\lambda \underline{h})}\right), \quad (50)$$

and upon choosing $q = (\varphi_x(\underline{h}\lambda)/n)^{-1/(2(\delta+1))}$ we have $o(\varphi^{-1}q^{-(\delta+1)}) = o(\varphi^{-1}(\varphi/n)^{1/2}) = o(n^{-1/2}\varphi^{-1/2}) = o(1)$, and also

$$\begin{aligned} O\left(\frac{1}{\varphi_x(\underline{h}\lambda)} q\sqrt{v_2(m_n)}\right) &= O\left(\frac{1}{\varphi_x(\underline{h}\lambda)} \cdot \left(\frac{\varphi_x(\underline{h}\lambda)}{n}\right)^{-1/(2(\delta+1))} \sqrt{v_2(m_n)}\right) \\ &= O\left(\frac{\sqrt{v_2(m_n)}}{[\varphi_x(\underline{h}\lambda)]^{(2\delta+3)/(2\delta+2)} n^{-1/(2(\delta+1))}}\right) = o(1) \end{aligned} \quad (51)$$

by Assumption S2. Hence $R_3 = o(1)$ and similarly $R_2 = o_p(1)$.

As for the first term that remains, we can rewrite R_1 as

$$\frac{1}{n} \sum_{t=1}^n \left[\frac{K_{t,(m)} Y_t - E(K_{t,(m)} Y_t)}{EK_{1,(m)}} \right] + \frac{1}{n} \sum_{t=1}^n \left[\frac{\{K_{t,(m)} Y_t - E(K_{t,(m)} Y_t)\} \cdot \{EK_{1,(m)} - EK_1\}}{EK_{1,(m)} \cdot EK_1} \right],$$

and denote by each summation as F_1 and F_2 , respectively.

On noting that $K_{t,(m)}$ is α -mixing, we follow the arguments we used for the static regression case. Write $(K_{t,(m)} Y_t - E(K_{t,(m)} Y_t))/EK_{1,(m)} = Q_{t,(m)}$; then since Q_t is near epoch dependent on Y_t thanks to (the extra factor φ^{-1} in) condition (13) stipulated on the rate of stability, by using the approximation scheme one can readily see that the first component $F_1 \rightarrow^p 0$.

Furthermore, moving on to the next term we see that

$$F_2 \leq \left[\frac{1}{n} \sum_{t=1}^n \frac{\{K_{t,(m)} Y_t - E(K_{t,(m)} Y_t)\}}{EK_{1,(m)}} \right] \cdot \frac{C\sqrt{v_2(m_n)}}{EK_1} = o_p\left(\frac{\sqrt{v_2(m_n)}}{\varphi_x(\underline{h}\lambda)}\right), \quad (52)$$

completing the proof. ■

4.2 Proof of Theorem 2 and 3

With additional sumptions A7, A8 and B3 it follows that

$$\begin{aligned} \mathcal{B}_n &= E\hat{m}_2(x) - m(x) = E\left(\frac{1}{nEK_1} \sum_{t=1}^n K_t Y_t - m(x)\right) \\ &= \frac{1}{EK_1} EK_1 Y_1 - \frac{EK_1}{EK_1} m(x) = \frac{1}{EK_1} E\left[E\left[(Y_1 - m(x))K_1 \middle| X\right]\right] \\ &= \frac{1}{EK_1} E\left[\left[m(X) - m(x)\right]K_1\right] \leq \sup_{u \in \mathcal{E}(x, \lambda \underline{h})} |m(u) - m(x)| \\ &\leq \sup_{u \in \mathcal{E}(x, \lambda \underline{h})} \sum_{j=1}^{\infty} c_j |u_j - x_j|^\beta = \sum_{j=1}^{\infty} c_j (\lambda h \phi_j)^\beta = h^\beta \left(\lambda^\beta \sum_{j=1}^{\infty} c_j j^{p\beta} \right) < \infty. \end{aligned} \quad (53)$$

Moving on, rewriting the decomposition (15) as

$$\widehat{m}(x) - m(x) - B(x) = \frac{B(x) \cdot [1 - \widehat{m}_1(x)]}{\widehat{m}_1(x)} + \frac{\widehat{m}_2(x) - E\widehat{m}_2(x) - m(x)[\widehat{m}_1(x) - 1]}{\widehat{m}_1(x)},$$

we see that it suffices to derive the limiting distribution of

$$\begin{aligned} & \widehat{m}_2(x) - E\widehat{m}_2(x) - m(x)[\widehat{m}_1(x) - 1] \\ &= \frac{1}{n} \sum_{t=1}^n \frac{1}{EK_1} [K_t Y_t - m(x)K_t - E(K_t Y_t) + m(x)EK_t] =: \frac{1}{n} \sum_{t=1}^n R_{nt}, \end{aligned} \quad (54)$$

since $\widehat{m}_1(x) \xrightarrow{p} 1$ as a consequence of Theorem 1. Now, with Assumption A6, B3, B4, and the law of iterated expectations, the asymptotic variance of (54) is given by

$$\begin{aligned} \frac{\text{Var}[K_t(Y_t - m(x))]}{nE^2K_1} &= \frac{1}{nE^2K_1} \left\{ E \left[K_t(Y_t - m(x)) \right]^2 - E^2[K_t(Y_t - m(x))] \right\} \\ &= \frac{1}{nE^2K_1} \left\{ E[\sigma^2(X)K_1^2] + E \left([m(X) - m(x)]^2 K_1^2 \right) \right\} \\ &= \frac{1}{nE^2K_1} \left\{ \sigma^2(x)EK_1^2 + E \left([\sigma^2(X) - \sigma^2(x)] K_1^2 \right) + o(1)EK_1^2 \right\} \\ &= \frac{EK_1^2}{nE^2K_1} (\sigma^2(x) + o(1)) \simeq \frac{\sigma^2(x)\xi_2}{n\varphi_x(\underline{h}\lambda)\xi_1^2}. \end{aligned} \quad (55)$$

Following similar arguments and using the latter assumption of B4, it can be readily shown that the covariance term is of smaller order than (55).

Noting (53) and (55) we consider the normalized statistic $R_{nt}^* := \sqrt{\varphi_x(\underline{h}\lambda)} \cdot R_{nt}$ and derive the limiting distribution of $(1/\sqrt{n}) \cdot R_{nt}^*$. The usual procedure for establishing the asymptotic normality for the case of dependent observations is to employ Bernstein's blocking method; alternating blocks of two different sizes (hereafter referred to as the “big” and “small” blocks) are chosen to partition the index set $t = 1, \dots, n$. The argument proceeds by showing that the “small blocks” can be appropriately selected in such a way that they can be omitted in large sample, and then asymptotic independence can be obtained over the “big blocks”. One may then apply the standard arguments involving the Lindeberg-Feller central limit theorem upon checking the Lindeberg conditions.

Below we shall only prove the autoregression case, where an additional step of mixing approximation is added to the standard regression case. The asymptotic normality for the regression case was established by Masry (2005) in a general context; we follow and briefly sketch his proof for the sake of completeness. Partition the index set $\{1, \dots, n\}$ by $2k (= 2k_n \rightarrow \infty)$ number of blocks of two different sizes that alternate (called the “big-blocks” and the “small-blocks”, respectively), along with a single block (the “last block”) that covers the remainder. The size of the alternating blocks is given by a_n and b_n respectively, where the one for the “big-blocks” a_n is set to dominate that for the “small-blocks” b_n in large sample, i.e. $b_n = o(a_n)$. More specifically, we take $k_n = \lfloor n/(a_n + b_n) \rfloor$ and $a_n = \lfloor \sqrt{n\varphi_x(\lambda\underline{h})}/q_n \rfloor$ where q_n is a sequence of integer that goes off to infinity; it then clearly follows that $a_n/n \rightarrow 0$ and $a_n/\sqrt{n\varphi_x(\lambda\underline{h})} \rightarrow 0$. We also assume $(n/a_n) \cdot \alpha^*(b_n) \rightarrow 0$, where α^* is the mixing coefficient for $R_{nt,(m)}^*$ below.

By construction above we can write $\sqrt{n}^{-1} \sum_{t=1}^n R_{nt}^*$ as the sum of the groups of big-blocks $\mathcal{B}$, small-blocks $\mathcal{S}$ and the remainder block $\mathcal{R}$ defined as

$$\begin{aligned}\mathcal{B} &:= \frac{1}{\sqrt{n}} \sum_{j=0}^{k-1} \Xi_{1,j} = \frac{1}{\sqrt{n}} \sum_{j=0}^{k-1} \left(\sum_{t=j(a+b)+1}^{j(a+b)+a} R_{nt}^* \right) \\ \mathcal{S} &:= \frac{1}{\sqrt{n}} \sum_{j=0}^{k-1} \Xi_{2,j} = \frac{1}{\sqrt{n}} \sum_{j=0}^{k-1} \left(\sum_{t=j(a+b)+a+1}^{(j+1)(a+b)} R_{nt}^* \right) \\ \mathcal{R} &:= \frac{1}{\sqrt{n}} \Xi_{3,j} = \frac{1}{\sqrt{n}} \left(\sum_{t=k(a+b)+1}^n R_{nt}^* \right).\end{aligned}$$

The aim is to show that the contributions from the small and the last remaining block are negligible, and that the big-blocks are asymptotically independent.

We first consider the big blocks $\mathcal{B}$, writing

$$\mathcal{B} = \sum_{j=0}^{k-1} \left(\sum_{t=j(a+b)+1}^{j(a+b)+a} R_{nt,(m)}^* \right) + \sum_{j=0}^{k-1} \left(\sum_{t=j(a+b)+1}^{j(a+b)+a} R_{nt,(m)}^* - \sum_{t=j(a+b)+1}^{j(a+b)+a} R_{nt}^* \right) = \mathcal{Q}_1 + \mathcal{Q}_2,$$

where m is as in S2 and $R_{nt,(m)}^* = E(R_{nt}^* | Y_t, \dots, Y_{t-m+1})$. As for the term $\mathcal{Q}_2$, write

$$\begin{aligned}\mathcal{Q}_2 &= \sum_{j=0}^{k-1} \sum_{t=j(a+b)+1}^{j(a+b)+a} \left(R_{nt,(m)}^* - R_{nt}^* \right) \\ &= \sum_{j=0}^{k-1} \sum_{t=j(a+b)+1}^{j(a+b)+a} \frac{1}{EK_1} \left[K_t Y_t - K_{t,(m)} Y_t - E(K_t Y_t) + E(K_{t,(m)} Y_t) \right] \\ &\quad - \sum_{j=0}^{k-1} \sum_{t=j(a+b)+1}^{j(a+b)+a} \frac{1}{EK_1} \cdot m(x) \left[K_t - K_{t,(m)} - EK_t + EK_{t,(m)} \right] = \mathcal{I}_1 + \mathcal{I}_2.\end{aligned}$$

Noting that the terms being summed up in $\mathcal{I}_1$ are the same as those in $R_2 + R_3$ in p. 24 and that R_{nt}^* is near epoch dependent on Y_t it can be easily seen that $\sqrt{n}^{-1} \mathcal{Q}_2 = o_p(1)$ by following similar arguments used before, and using the conditions on a_n, b_n and k_n . Further, the asymptotic independence of terms in $\mathcal{Q}_1$ holds by the Volkonskii-Rozanov inequality (see Fan and Yao (2003, p. 72)), and by the fact that $(n/a_n) \cdot \alpha^*(b_n) \rightarrow 0$.

Moving on to the small blocks, due to stationarity we have

$$\begin{aligned}\text{Var}(\mathcal{S}) &= \frac{1}{n} \text{Var} \left(\sum_{j=0}^{k-1} \sum_{t=j(a+b)+a+1}^{(j+1)(a+b)} R_{nt}^* \right) \\ &= \frac{1}{n} \sum_{j=0}^{k-1} \text{Var} \left(\sum_{t=j(a+b)+a+1}^{(j+1)(a+b)} R_{nt}^* \right) + \frac{1}{n} \sum_{j \neq l}^{k-1} \text{Cov} \left(\sum_{t=j(a+b)+a+1}^{(j+1)(a+b)} R_{nt}^*, \sum_{s=l(a+b)+a+1}^{(l+1)(a+b)} R_{ns}^* \right) \\ &= \frac{1}{n} \sum_{j=0}^{k-1} \left(b_n \text{Var}(R_{nt}^*) + \sum_{t \neq l}^{b_n} \text{Cov}(R_{nt}^*, R_{nl}^*) \right) + \frac{1}{n} \sum_{j \neq l}^{k-1} \sum_{i,j=1}^{b_n} \text{Cov}(R_{n,i+w_j}^*, R_{n,r+w_l}^*) \\ &= \mathcal{Q}_1 + \mathcal{Q}_2 + \mathcal{Q}_3.\end{aligned}$$

where $w_j = j(a + b) + a$.

Regarding the first term, similar arguments used in deriving (55) yield

$$Q_1 = \frac{1}{n} k_n b_n \frac{[\varphi_x(\underline{h}\lambda)^{1/2}]^2 \sigma^2(x) \xi_2}{\varphi_x(\underline{h}\lambda) \xi_1^2} = \frac{k_n b_n \sigma^2(x) \xi_2}{n \xi_1^2} \rightarrow 0 \quad (56)$$

because $k_n b_n / n \sim b_n / (a_n + b_n) \rightarrow 0$. Now moving on to Q_2 and Q_3 , the sum of covariances can be dealt with in the same manner as we did for the variance using (55), so $Q_2 \rightarrow 0$. Similarly for Q_3 , implying $\text{Var}(\mathcal{S}) \rightarrow 0$ as desired. Convergence result for the remainder $\mathcal{R}$ can be established similarly, and is bounded by $C(a_n + b_n)/n \rightarrow 0$.

Consequently, upon checking the Lindeberg conditions it follows that

$$\frac{\sum_t R_{nt} - E R_{nt}}{\sqrt{n \varphi_x(\lambda \underline{h})^{-1} \sigma^2(x) \xi_2 \xi_1^{-2}}} = \frac{m_2(x) - E \hat{m}_2(x) - m(x) [\hat{m}_1(x) - 1]}{\left(\sqrt{n \varphi_x(\lambda \underline{h})} \right)^{-1} \sqrt{\sigma^2(x) \xi_2 \xi_1^{-2}}} \Rightarrow N(0, 1). \quad (57)$$

The rest of the proof concentrates on specification of the small ball probability:

$$\begin{aligned} \varphi_x(\lambda \underline{h}) &= P(\|H^{-1}(x - X)\| \leq 1 \cdot \lambda) = P(X \in \mathcal{E}(x, \lambda \underline{h})) \\ &= P\left(\sum_{j=1}^{\infty} j^{-2p} (x_j - X_j)^2 \leq h^2 \lambda^2\right) = P(\|z - Z\| \leq h \lambda) \\ &= \int_{B(0, h \lambda)} dP_{z-Z}(u) = \int_{B(0, h \lambda)} p^*(u) dP_Z(u) \simeq p^*(0) \cdot P(\|Z\| \leq h \lambda) \end{aligned} \quad (58)$$

for any constant vector $z \in \ell_2$, where $p^*(\cdot)$ is the Radon-Nikodym derivative of the induced probability measure P_{z-Z} with respect to P_Z (cf. Assumption B1); the explicit form of the derivative is not known in general. Nonetheless, in the special case of Gaussian process it is well-known that

$$P(\|z - Z\| \leq \epsilon) \simeq P(\|Z\| \leq \epsilon) \exp\left\{-\frac{1}{2} \|\Gamma^{-1/2} z\|^2\right\} \quad \text{as } \epsilon \rightarrow 0, \quad (59)$$

where the asymptotic equivalence is obtained by establishing the upper and lower bounds using Anderson's inequality (e.g. Proposition 2, Lewandowski et al. (1995)) and the Cameron-Martin theorem, respectively. See for instance Zolotarev (1988), or Li and Shao (2001) for detailed discussion.

Having reduced the problem of shifted small deviation probability to the non-shifted case, we can now specify the explicit rate of convergence upon substituting $r = h^2 \lambda^2$, $A = 2p$, and $a = 2p/(2p - 1)$ in Proposition 4.1 of Dunker et al. (1998) for the i.i.d. case. In the dependent case, i.e. when the regressors satisfy Assumption D2, the small ball probability can be specified in view of Theorem 1 of Hong, Lifshits and Nazarov (2016), completing the proof. $\blacksquare$

4.3 Proof of Lemma 1 and 2

Lemma 1 is a straightforward extension of Lemma 4.3 and 4.4 of Ferraty and Vieu (2006), and hence is omitted. Lemma 2 can be shown by noting that for each n the τ_n -dimensional polyhedron $D := \{w = (w_i)_{i \leq \tau} \in \mathbb{R}^\tau, |w_i| \leq \lambda\}$ can be covered by $(\lceil 2\lambda\sqrt{\tau}/\varepsilon \rceil + 1)^\tau$ number of balls of radius ε , see Chaté and Courbage (1997), and then following the proof of Theorem 2 in Jia et al. (2003). $\blacksquare$

4.4 Proof of Theorem 4

As before, we start from the decomposition (15):

$$\widehat{m}(x) - m(x) = \frac{1}{\widehat{m}_1(x)} \left(\left[\widehat{m}_2(x) - E\widehat{m}_2(x) \right] + \left[E\widehat{m}_2(x) - m(x) \right] - m(x) \left[\widehat{m}_1(x) - 1 \right] \right).$$

Noting that the small deviation for $X = (X_1, \dots, X_\tau, 0, 0, \dots)$ is

$$\varphi_x(\lambda \underline{h}) = P \left(\sum_{j=1}^{\tau} j^{-2p} X_j^2 \leq h^2 \right) \simeq P \left(\sum_{j=1}^{\infty} j^{-2p} X_j^2 \leq h^2 + \tau^{-2p+1} \right),$$

the proof reduces to showing three steps, where in the first step we show

$$\sup_{x \in \mathcal{S}_\tau} \left| \widehat{m}_2(x) - E\widehat{m}_2(x) \right| = O_{a.s.} \left(\sqrt{\frac{(\log n)^2}{n \varphi_x(\lambda \underline{h})}} \right). \quad (60)$$

We cover the set $\mathcal{S}_\tau$ defined in (30) by $L = L(\mathcal{S}_\tau, \eta)$ number of balls of radius η denoted I_k , each of which is centred at x_k , for $k = 1, \dots, L$. i.e. $\mathcal{S}_\tau \subset \bigcup_{k=1}^{L_n} B(x_{k,n}, \eta)$. Then it follows that

$$\begin{aligned} \sup_{x \in \mathcal{S}_\tau} \left| \widehat{m}_2(x) - E\widehat{m}_2(x) \right| &= \max_{1 \leq k \leq L_n} \sup_{x \in I_k \cap \mathcal{S}_\tau} \left| \widehat{m}_2(x) - E\widehat{m}_2(x) \right| \\ &= \max_{1 \leq k \leq L_n} \sup_{x \in I_k \cap \mathcal{S}_\tau} \left| \widehat{m}_2(x) - \widehat{m}_2(x_k) + \widehat{m}_2(x_k) - E\widehat{m}_2(x_k) + E\widehat{m}_2(x_k) - E\widehat{m}_2(x) \right| \\ &\leq \max_{1 \leq k \leq L_n} \sup_{x \in I_k \cap \mathcal{S}_\tau} \left| \widehat{m}_2(x) - \widehat{m}_2(x_k) \right| + \max_{1 \leq k \leq L_n} \sup_{x \in I_k \cap \mathcal{S}_\tau} \left| E\widehat{m}_2(x_k) - E\widehat{m}_2(x) \right| \\ &\quad + \max_{1 \leq k \leq L_n} \left| \widehat{m}_2(x_k) - E\widehat{m}_2(x_k) \right| =: R_1 + R_2 + R_3. \end{aligned} \quad (61)$$

We first consider R_1 :

$$\begin{aligned} R_1 &= \max_{1 \leq k \leq L_n} \sup_{x \in I_k \cap \mathcal{S}_\tau} \left| \widehat{m}_2(x) - \widehat{m}_2(x_k) \right| \\ &= \max_{1 \leq k \leq L_n} \sup_{x \in I_k \cap \mathcal{S}_\tau} \left| \frac{1}{nEK_1} \sum_{t=1}^n Y_t K \left(\|H^{-1}(x - X_t)\| \right) - Y_t K \left(\|H^{-1}(x_k - X_t)\| \right) \right| \\ &\leq \max_{1 \leq k \leq L_n} \sup_{x \in I_k \cap \mathcal{S}_\tau} \frac{C}{n \varphi_x(\lambda \underline{h})} \sum_{t=1}^n |Y_t K_t - Y_t K_{t,k}| \cdot \mathbf{1}_{\mathcal{E}(x, \lambda \underline{h})}(X_t) \cdot \mathbf{1}_{\mathcal{E}(x_k, \lambda \underline{h})}(X_t), \end{aligned}$$

where $K_{t,k} := K(\|H^{-1}(x_k - X_t)\|)$. Now, because the kernel function is assumed to be Lipschitz continuous by Assumption C2, on choosing $\eta = \log n/n$ we get

$$R_1 \leq \frac{1}{n} \sum_{t=1}^n \frac{|Y_t|}{\varphi_x(\underline{h} \lambda)} \eta h^{-1} \mathbf{1}_{\mathcal{E}(x, \lambda \underline{h}) \cup \mathcal{E}(x_k, \lambda \underline{h})}(X_t) =: \frac{1}{n} \sum_{t=1}^n J_t,$$

where J_t is α -mixing under both assumptions S1' and S2'. Then for some $\delta > 0$

$$E|J_t|^{2+\delta} \leq \frac{E|Y_t|^{2+\delta} \eta^{2+\delta}}{\varphi_x(\lambda \underline{h})^{2+\delta} h^{2+\delta}} \varphi_x(\lambda \underline{h}) \leq C \left(\frac{\log n}{n \varphi_x(\lambda \underline{h}) h} \right)^{1+\delta} \times \left(\frac{\log n}{nh} \right) \leq C' \quad (62)$$

by Assumption A4 and C1, which suggests that for any positive t

$$P(|J_t| > t) \leq \frac{E|J_t|^{2+\delta}}{t^{2+\delta}} \leq C't^{-(2+\delta)}. \quad (63)$$

Also, by Assumption A6 we have

$$E|J_t| \leq \frac{E(E(|Y_t||X))\eta}{\varphi_x(\lambda\underline{h})h} \varphi_x(\lambda\underline{h}) \leq \frac{C\eta}{h}. \quad (64)$$

By Lemma 2 we can specify the Kolmogorov's entropy for S_τ with $\eta = \log n/n$:

$$\log L\left(S, \frac{\log n}{n}\right) = C \log \left[\left(\frac{2\lambda n}{\sqrt{\log n}} + 1 \right)^{\log n} \right] \sim \log n \times \log \left[\frac{2\lambda n}{\sqrt{\log n}} \right]$$

for sufficiently large n and λ , implying that the order of Kolmogorov's $\frac{\log n}{n}$ entropy is

$$O\left(\log L\left(S_\tau, \frac{\log n}{n}\right)\right) = O\left((\log n)^2 - \log n[\log \log n]\right) = O\left((\log n)^2\right). \quad (65)$$

Now, with (63) we may apply the Fuk-Nagaev inequality⁵ with $r = (\log n)^2$ and $\varepsilon = \varepsilon_0[\log L(S, \frac{\log n}{n})/(n\varphi(\lambda\underline{h}))]^{1/2}$ for some positive constant ε_0 . Then since

$$s_n^2 := \sum_{t=1}^n \sum_{s=1}^n \text{Cov}(J_t, J_s) \leq C \left(\frac{1}{\varphi(\lambda\underline{h})} \frac{(\log n)^2}{n^2 h^2} - \frac{(\log n)^2}{n^2 h^2} \right) = O\left(\frac{n \log n}{\varphi(\lambda\underline{h})}\right),$$

and because an exponentially α -mixing sequence is arithmetically α -mixing with some mixing rate $k > 0$, it follows that

$$\begin{aligned} P\left(\frac{1}{n} \left| \sum_{t=1}^n J_t - EJ_1 \right| > \varepsilon_0 \sqrt{\frac{\log L(S, \frac{\log n}{n})}{n\varphi(\lambda\underline{h})}}\right) \\ \leq 4 \left(1 + \frac{n^2 \varepsilon_0^2 \log L(S, \frac{\log n}{n})}{16(\log n)^2 s_n^2 n \varphi(\lambda\underline{h})} \right)^{-\frac{(\log n)^2}{2}} + \frac{C' \cdot n}{(\log n)^2} \left(\frac{(\log n)^4 n \varphi(\lambda\underline{h})}{n^2 \varepsilon_0^2 \log L(S, \frac{\log n}{n})} \right)^{\frac{(k+1)(2+\delta)}{2(2+\delta+k)}} \\ = 4 \left(1 + \frac{n}{16 s_n^2 \varphi(\lambda\underline{h}) \varepsilon_0^2} \right)^{-\frac{(\log n)^2}{2}} + \frac{C' \cdot n}{(\log n)^2} \left(\frac{\varphi(\lambda\underline{h})}{n \varepsilon_0^2} \right)^{\frac{2k+k\delta+2+\delta}{4+2\delta+2k}} (\log n)^{\frac{(k+1)(2+\delta)}{2+\delta+k}} \\ \leq 4 \exp\left(-\frac{\varepsilon_0^2 (\log n)^2 n}{32 s_n^2 \varphi(\lambda\underline{h})}\right) + C' n^{-\frac{k\delta-(\delta+2)}{2(2+\delta+k)}} (\log n)^{\frac{k\delta-(\delta+2)}{2+\delta+k}} \varphi(\lambda\underline{h})^{\frac{2k+k\delta+2+\delta}{2(2+\delta+k)}} \\ \leq 4 \exp\left(-\frac{\varepsilon_0^2 \log n}{32}\right) + C' \left(\frac{\log n}{n^2}\right)^{\frac{k\delta-(\delta+2)}{2+\delta+k}} \end{aligned} \quad (66)$$

by choosing k sufficiently large so that $k\delta > \delta + 2$ and Taylor expansion of $\log(1 + \epsilon)$ for sufficiently small $\epsilon > 0$. Now summing up over $n = 1$ upto infinity, we see that

$$\begin{aligned} \sum_{n=1}^{\infty} P\left(\frac{1}{n} \left| \sum_{i=1}^n J_i - EJ_1 \right| > \varepsilon_0 \sqrt{\frac{(\log n)^2}{n\varphi(\lambda\underline{h})}}\right) \\ \leq C \sum_{n=1}^{\infty} \exp\left(-\frac{\varepsilon_0^2 \log n}{32}\right) + \sum_{n=1}^{\infty} \left(\frac{\log n}{n^2}\right)^{\frac{k\delta-(\delta+2)}{2+\delta+k}} < \infty \end{aligned}$$

⁵See Rio (2000, p. 87) for details; the exact bounds of the constants are specified.

by choosing ε_0 sufficiently big (i.e. any finite real number strictly bigger than $4\sqrt{2}$).

Hence by (64) and by the Borel-Cantelli lemma we have

$$\begin{aligned} R_1 &= \max_{1 \leq k \leq L_n} \sup_{x \in I_k \cap \mathcal{S}_\tau} |\hat{m}_2(x) - \hat{m}_2(x_k)| \leq O\left(\frac{\eta}{h}\right) + O_{a.s.}\left(\sqrt{\frac{(\log n)^2}{n\varphi_x(\lambda \underline{h})}}\right) \\ &= O\left(\sqrt{\frac{(\log n)^2}{n^2 h^2}}\right) + O_{a.s.}\left(\sqrt{\frac{(\log n)^2}{n\varphi_x(\lambda \underline{h})}}\right) = O_{a.s.}\left(\sqrt{\frac{(\log n)^2}{n\varphi_x(\lambda \underline{h})}}\right). \end{aligned} \quad (67)$$

For the second term R_2 , following similar arguments as above we can show that

$$R_2 \leq \max_{1 \leq k \leq L_n} \sup_{x \in I_k \cap \mathcal{S}_\tau} E|\hat{m}_2(x) - \hat{m}_2(x_k)| = O\left(\sqrt{\frac{(\log n)^2}{n\varphi_x(\lambda \underline{h})}}\right). \quad (68)$$

Next we study the last component:

$$R_3 = \max_{1 \leq k \leq L_n} |\hat{m}_2(x_k) - E\hat{m}_2(x_k)| =: \max_{1 \leq k \leq L_n} |W_n(x_k)| \quad (69)$$

where

$$\begin{aligned} W_n(x) &= \hat{m}_2(x) - E\hat{m}_2(x) = \frac{1}{nEK_1} \sum_{t=1}^n [Y_t K_t - EY_t K_t] \\ &= \frac{1}{n} \sum_{t=1}^n \left[\frac{Y_t K_t}{EK_1} - \frac{EY_t K_t}{EK_1} \right] =: \frac{1}{n} \sum_{t=1}^n Q_t \end{aligned}$$

as defined in (37), and by elementary arguments

$$P\left(\max_{1 \leq k \leq L_n} |\hat{m}_2(x_k) - E\hat{m}_2(x_k)| > \varepsilon\right) \leq L_n \cdot \sup_{x \in \mathcal{S}} P(|W_n(x)| > \varepsilon). \quad (70)$$

As we did before in the pointwise case, we must separately consider the cases of regression and autoregression because the asymptotic arguments to follow depends upon the temporal dependence structure of Q_t .

In the regression case, we use the exponential inequality of Bosq (1996, Theorem 1.3.2) for α -mixing sequence. We note that $|Q_t| \leq C/\varphi_x(\underline{h}\lambda) =: b$ for all t , and that $\sigma^2(m) := p \cdot \text{Var}(Q_t) = O(p/\varphi(\underline{h}\lambda))$ (where $p = n/(2q)$ and $q = \log n \sqrt{n}/\sqrt{\varphi}$) by the Cauchy-Schwarz inequality and assumption A4. Hence, it follows that

$$v^2(m) = \frac{2}{p^2} \sigma^2(m) + \frac{b\varepsilon}{2} \leq \frac{Cq}{n\varphi_x(\underline{h}\lambda)} + \frac{C\varepsilon}{\varphi_x(\underline{h}\lambda)} \leq \frac{C'\varepsilon}{\varphi_x(\underline{h}\lambda)},$$

where $\varepsilon = \varepsilon_0 \sqrt{\log L_n/(n\varphi)}$, and by assumption S1 that

$$\begin{aligned} P\left(\left|\hat{m}_2(x) - E\hat{m}_2(x)\right| > \varepsilon_0 \sqrt{\frac{\log L(S, \frac{\log n}{n})}{n\varphi(\lambda \underline{h})}}\right) \\ \leq 4 \exp\left\{-\frac{\varepsilon^2 q}{8v^2(m)}\right\} + 22\left(1 + \frac{4b}{\varepsilon}\right)^{1/2} q\alpha\left(\left[\frac{n}{2q}\right]\right) \\ \leq 4 \exp\left\{-\frac{\varepsilon_0 q \varphi \sqrt{\log L_n}}{8\sqrt{n\varphi}}\right\} + 22\left(1 + \frac{4\sqrt{n\varphi}}{\varphi \log n}\right)^{1/2} \frac{\log n \sqrt{n}}{\sqrt{\varphi}} \alpha\left(\left[\frac{\sqrt{n\varphi}}{2 \log n}\right]\right) \\ \leq 4 \exp\left\{-\frac{\varepsilon_0 \log L_n}{8}\right\} \leq CL_n^{-\varepsilon_0/8}, \end{aligned} \quad (71)$$

due to exponential decay of $\alpha(\cdot)$; with (70) the desired result follows by choosing ε_0 large enough.

In the autoregression case (i.e. under assumption D2), using the mixing approximation techniques adopted in Section 4.1. and following the arguments in (46), (51) and (52), it can be readily shown that the same conclusion can be derived; that is:

$$R_3 = \max_{1 \leq k \leq L_n} \left| \widehat{m}_2(x_k) - E\widehat{m}_2(x_k) \right| = O_{a.s.} \left(\sqrt{\frac{(\log n)^2}{n\varphi_x(\lambda \underline{h})}} \right). \quad (72)$$

Now returning back to where we started, viewing $\widehat{m}_1(x)$ as a special case of $\widehat{m}_2(x)$ with $Y_t = 1 \ \forall t$, we can repeat the above procedure, yielding (since $E\widehat{m}_1(x) = 1$)

$$\sup_{x \in \mathcal{S}_\tau} \left| \widehat{m}_1(x) - 1 \right| = O_{a.s.} \left(\sqrt{\frac{(\log n)^2}{n\varphi_x(\lambda \underline{h})}} \right). \quad (73)$$

Finally the proof is complete in view of (60), (67), (68), (72), (73), and the contribution from the bias component. ■

References

- [1] de Acosta, A., 1983. *Small deviations in the functional central limit theorem with applications to functional laws of the iterated logarithm*. Annals of Probability, 11(1), 78-101.
- [2] Andrews, D. W., 1995. *Nonparametric kernel estimation for semiparametric models*. Econometric Theory, 11(3), 560-586.
- [3] Bierens, H. J., 1983. *Uniform consistency of kernel estimators of a regression function under generalized conditions*. Journal of the American Statistical Association, 78(383), 699-707.
- [4] Bierens, H. J., 1987. *Kernel estimators of regression functions*. In Advances in econometrics: Fifth world congress, 1, 99-144.
- [5] Bingham, N. H., Goldie, C. M., Teugels, J. L., 1987. *Regular Variation*. Cambridge University Press.
- [6] Bochner, S., 1955. *Harmonic Analysis and the Theory of Probability*. University of California Press, Berkeley and Los Angeles.
- [7] Bosq, B., 1996. *Nonparametric statistics for stochastic processes: estimation and prediction*. Springer-Verlag, New York.
- [8] Bradley, R. C., 2005. *Basic properties of strong mixing conditions. A survey and some open questions*. Probability Surveys, 2(2), 107-144.
- [9] Bulinskii, A. V., Shiryaev, A. N., 2003. *Theory of Random Processes* (in Russian). Fizmatlit.

- [10] Chaté, H., Courbage, M. (ed.), 1997. *Special issue on lattice dynamics*. Physica D, 103, 1-612.
- [11] Davidson, J., 1994. *Stochastic Limit Theory: An Introduction for Econometricians: An Introduction for Econometricians*. Oxford University Press.
- [12] Davydov, Y. A., 1968. *Convergence of distributions generated by stationary stochastic processes*. Theory of Probability and Applications, 13(4), 691-696.
- [13] Delsol, L., 2009. *Advances on asymptotic normality in non-parametric functional time series analysis*. Statistics, 43(1), 13-33.
- [14] Devroye, L. P., 1978. *The uniform convergence of the Nadaraya-Watson regression function estimate*. Canadian Journal of Statistics, 6(2), 179-191.
- [15] Devroye, L., 1981. *On the almost everywhere convergence of nonparametric regression function estimates*. Annals of Statistics, 9(6), 1310-1319.
- [16] Doukhan, P., 1994. *Mixing*. Springer, New York.
- [17] Doukhan, P., Wintenberger, O., 2008. *Weakly dependent chains with infinite memory*. Stochastic Processes and their Applications, 118(11), 1997-2013.
- [18] Duflo, M., 1997. *Random iterative models*. Springer-Verlag, Berlin.
- [19] Dunker, T., Lifshits, M. A., Linde, W., 1998. *Small deviation probabilities of sums of independent random variables*. In: Eberlein, E. (ed.), High dimensional probability, volume 43 of Progress in Probability., Birkhauser, Basel, 59-74.
- [20] Fan, J., Masry, E., 1992. *Multivariate regression estimation with errors-in-variables: asymptotic normality for mixing processes*. Journal of Multivariate Analysis, 43(2), 237-271.
- [21] Fan, J., 1990. *A remedy to regression estimators and nonparametric minimax efficiency*. Department of Statistics, University of North Carolina at Chapel Hill.
- [22] Fan, J., Yao, Q., 2005. *Nonlinear Time Series: Nonparametric and Parametric Methods*. Springer, New York.
- [23] Feller, W., 1971. *Introduction to Probability Theory and Its Applications*, Vol. 2. Wiley, New York.
- [24] Ferraty, F., Laksaci, A., Tadj, A., Vieu, P., 2010. *Rate of uniform consistency for nonparametric estimates with functional variables*. Journal of Statistical Planning and Inference, 140, 335-352.
- [25] Ferraty, F., Romain, Y., 2010. *The Oxford Handbook of Functional Data Analysis*. Oxford University Press.
- [26] Ferraty, F., Vieu, P., 2002. *The functional nonparametric model and application to spectrometric data*. Computational Statistics, 17, 545-564.

- [27] Ferraty, F., Vieu, P., 2006. *Nonparametric functional data analysis: Theory and Practice*. Springer, New York.
- [28] Fuk, D. K., Nagaev, S. V., 1971. *Probability inequalities for sums of independent random variables*. Theory of Probability and its applications, 16, 643-660.
- [29] Götze, F., Hipp, C., 1994. *Asymptotic distribution of statistics in time series*. Annals of Statistics, 22(4), 2062-2088.
- [30] Greblicki, W., Krzyzak, A., 1980. *Asymptotic properties of kernel estimates of a regression function*. Journal of Statistical Planning and Inference, 4(1), 81-90.
- [31] Geenens, G., 2011. *Curse of dimensionality and related issues in nonparametric functional regression*. Statistics Surveys, 5, 30-43.
- [32] Härdle, W. K., 1990. *Applied Nonparametric Regression*. (Vol. 27). Cambridge University Press.
- [33] Hong, S. Y., Lifshits, M., Nazarov, A., 2015. *Small Deviations for Dependent Sequences*. Preprint available at: <http://arxiv.org/abs/1511.05370>.
- [34] Ibragimov, I.A., Linnik, Y. B., 1971. *Independent and Stationary Sequences of Random Variables*. Wolters-Noordhoff, Groningen.
- [35] Jia, Q., Zhou, S., Yin, F., 2003. *Kolmogorov entropy of global attractor for dissipative lattice dynamical systems*. Journal of Mathematical Physics, 44, 5804-5801.
- [36] Karamata, J., 1933. *Sur un mode de croissance reguliere. Theoremes fondamentaux* (in French). Bulletin de la Societe Mathematique de France, 61, 55-62.
- [37] Lewandowski, M., Ryznar, M., 1995. *Anderson inequality is strict for Gaussian and stable measures*. Proceedings of the American Mathematical Society, 123(12), 3875-3880.
- [38] Li, W. V., Shao, Q. M., 2001. *Gaussian processes: inequalities, small ball probabilities and applications*. Handbook of Statistics, 19, 533-597.
- [39] Lifshits, M. A., 1997. *On the lower tail probabilities of some random series*. Annals of Probability, 25(1), 424-442.
- [40] Linton, O. B., Sancetta, A., 2009. *Consistent estimation of a general nonparametric regression function in time series*. Journal of Econometrics, 152, 70-78.
- [41] Lu, Z., 2001. *Asymptotic normality of kernel density estimators under dependence*. Annals of the Institute of Statistical Mathematics, 53(3), 447-468.
- [42] Mas, A., 2012. *Lower bound in regression for functional data by small ball probability representation in Hilbert space*, Electronic Journal of Statistics, 6, 1745-1778.
- [43] Masry, E., 2005. *Nonparametric regression estimation for dependent functional data: asymptotic normality*. Stochastic Processes and their Applications, 115, 155-177.

- [44] Nadaraya, E. A., 1964. *On estimating regression*. Theory of Probability & Its Applications, 9(1), 141-142.
- [45] Nadaraya, E. A., 1970. *Remarks on non-parametric estimates for density functions and regression curves*. Theory of Probability & Its Applications, 15(1), 134-137.
- [46] Pagan, A., Ullah, A., 1988. *The econometric analysis of models with risk terms*. Journal of Applied Econometrics, 3(2), 87-105.
- [47] Parzen, E., 1962. *On estimation of a probability density function and mode*. Annals of Mathematical Statistics, 33(3), 1065-1076.
- [48] Phillips, P. C., Park, J. Y., 1998. *Nonstationary density estimation and kernel autoregression*. Cowles Foundation for Research in Economics.
- [49] Ramsay, J. O., Silverman, B. W., 2002. *Applied functional data analysis: methods and case studies*. Springer, New York.
- [50] Rio, E., 2000. *Theorie asymptotique des processus aléatoires faiblement dépendants* (French) Springer, Mathématiques and Applications, 31.
- [51] Robinson, P. M., 1983. *Nonparametric estimators for time series*. Journal of Time Series Analysis, 4(3), 185-207.
- [52] Rosenblatt, M., 1956. *A central limit theorem and a strong mixing condition*. Proceedings of the National Academy of Sciences, 42(1), 43-47.
- [53] Roussas, G. G., 1989. *Consistent regression estimation with fixed design points under dependence conditions*. Statistics & Probability Letters, 8(1), 41-50.
- [54] Roussas, G. G., 1990. *Nonparametric regression estimation under mixing conditions*. Stochastic Processes and Their Applications, 36(1), 107-116.
- [55] Schuster, E. F., 1972. *Joint asymptotic distribution of the estimated regression function at a finite number of distinct points*. Annals of Mathematical Statistics, 43(1), 84-88.
- [56] Skorohod, A. B., 1967. *On the densities of probability measures in functional spaces*. Proc. Fifth Berkeley Sympos. Math. Statist. and Probability (Berkeley, Calif., 1965/66) Vol. II: Contributions to Probability Theory, 1, 163-182. University of California Press, Berkeley, CA.
- [57] Stone, C., 1980. *Optimal rates of convergence for nonparametric estimators*. Annals of Statistics, 8(6), 1348-1360.
- [58] Stone, C., 1982. *Optimal global rates of convergence for nonparametric regression*. Annals of Statistics, 10(4), 1040-1053.
- [59] Sytaya, G. N., 1974. *Certain asymptotic representations for a Gaussian measure in Hilbert space*. Theory of Random Process, 2, 93-104 (in Russian).
- [60] Watson, G. S., 1964. *Smooth regression analysis*, Sankhya Ser. A, 26, 359-372.

- [61] Wu, W. B., 2011. *Asymptotic theory for stationary processes*. Statistics and Its Interface, 0, 1-20.
- [62] Zolotarev, V. M., 1988. *Asymptotic behavior of the Gaussian measure in ℓ_2* . Journal of Soviet Mathematics, 24, 2330-2334.